

Figure E-5a. PSAT source categories by PM species contributions to the average 2000-2004 Baseline extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5b. PSAT source category by PM species contributions to the average 2018 projected extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5c. PSAT source region by source category contributions to the average 2000-2004 Baseline extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5d. PSAT source region by source category contributions to the average 2018 extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5e. Ranked PSAT source region by source category contributions to the average 2018 extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5f. Ranked PSAT source region by source category contributions to the average 2018 SO4 (left) and NO3 (right) extinction (Mm^{-1}) for the Worst 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5g. PSAT contributions by source category and PM species to the average 2000-2004 Baseline extinction (Mm^{-1}) for the Best 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5h. PSAT contributions by source category and PM species to the average 2018 extinction (Mm^{-1}) for the Best 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5i. PSAT contributions by source region and source category to the average 2000-2004 Baseline extinction (Mm^{-1}) for the Best 20% visibility days at Voyageurs (VOYA), Minnesota.




Figure E-5j. PSAT contributions by source region and source category to the average 2018 extinction (Mm^{-1}) for the Best 20% visibility days at Voyageurs (VOYA), Minnesota.