

Missouri Clean Water Commission Meeting  
Department of Natural Resources  
Lewis and Clark State Office Building  
LaCharrette/Nightingale Creek Conference Rooms  
1101 Riverside Drive  
Jefferson City, Missouri

January 06, 2016

**Public Hearing on the Draft 2018 303(d) Listing Methodology Document (LMD)**

**Issue:** The Federal Water Pollution Control Act, Section 303(d) requires states to biennially (once every two years) submit to the U.S. Environmental Protection Agency (EPA) a list of impaired waters for which adequate pollution controls have not yet been required.

**Background:** The department has a public participation process for revision of the Listing Methodology Document (LMD) that runs concurrently with the public notice for the 303(d) List. The proposed 2018 LMD was placed on public notice October 1, 2015 and will continue through January 31, 2016. All comments provided during the public availability meetings held on November 3, 2015 and December 1, 2015, and during the biological workgroup meeting held on November 18, 2015, can be found in the meeting discussion summaries provided in the Commission packet.

The draft 2016 LMD updates the 2014 LMD approved by the Commission in July 2014 and incorporates revisions related to reformatting and consolidation of information presented in the document, the addition of clarifying statements or information relating to biological assessment, and minor corrections to tables.

As of December 11, 2015 the department has received one (1) written comment on the proposed listing methodology document. The comment was received from the EPA Region 7. Written comments will continue to be accepted through January 31, 2016. All public comments, along with the department's responses, will become part of the administrative record for the LMD and will be made available on the department's website.

**Recommended Action:** No action is requested. This is an opportunity for staff, and the public, to present and comment on the draft 2018 Listing Methodology Document.

**Suggested Motion Language:** Hearing only.

**List of Attachments:**

- Proposed 2018 303(d) Listing Methodology Document. Additions from the 2016 LMD are shown in track changes and comment boxes.
- Summaries of Public Availability Meeting discussions held on November 3, 2015 and December 1, 2015 (under tab 1).
- Summary of Public Availability and Biological Workgroup Meeting discussions on the draft 2018 LMD held on November 18, 2015.

**Methodology for the Development  
of the  
2018 Section 303(d) List in Missouri**

Missouri Department of Natural Resources  
Division of Environmental Quality  
Water Protection Program



**Table of Contents**

**Comment [R1]:** Updated as needed

I. Citation and Requirements..... 1

    A. Citation of Section of Clean Water Act ..... 1

    B. U.S. EPA Guidance..... 1

II. The Methodology Document..... 7

    A. Procedures and Methods Used to Collect Water Quality Data..... 7

        • Department Monitoring ..... 7

        • Coordination with Other Monitoring Efforts in Missouri..... 7

        • Existing Monitoring Networks and Programs ..... 7

        • Identification of All Existing and Readily Available Water Quality Data Sources 12

        • Data Solicitation Request..... 12

        • Laboratory Analytical Support ..... 13

    B. Sources of Water Quality Data ..... 13

    C. Data Quality Considerations ..... 15

        • DNR Quality Assurance/Quality Control Program ..... 15

        • Other Quality Assurance/Quality Control Programs ..... 16

        • Other Data Quality Considerations ..... 16

    D. How Water Quality Data is Evaluated to Determine Whether or Not Waters are Impaired for 303(d) Listing Purposes ..... 18

        • Physical, Chemical, Biological and Toxicity Data ..... 18

        • Weight of Evidence Approach..... 18

        • Biological Data ..... 19

        • Other Biological Data ..... 31

        • Toxic Chemicals ..... 32

        • Duration of Assessment Period..... 33

        • Assessment of Tier Three Waters..... 34

        • Other Types of Information ..... 34

    E. Other 303(d) Listing Considerations ..... 35

    F. Prioritization of Waters for TMDL Development ..... 36

    G. Statistical Considerations..... 36

        • Description of Analytical Tools..... 37

        • Rationale for the Burden-of-Proof..... 37

        • Level of Significance Used in Tests ..... 38

        • Use of the Binomial Probability Distribution for Interpretation of the 10 Percent Rule..... 38

        • Other Statistical Considerations..... 39

        • Examples of Statistical Procedures..... 39

<u>H. References.....</u>	<u>41</u>
<u>Appendix A .....</u>	<u>43</u>
<u>Appendix B .....</u>	<u>46</u>
<u>Appendix C .....</u>	<u>50</u>
<u>Appendix D (11" x 14" Fold Out).....</u>	<u>55</u>

Proposed

## I. Citation and Requirements

### A. Citation of Section of Clean Water Act

The Missouri Department of Natural Resources (MDNR) is responsible for the implementation and administration of the Federal Clean Water Act in Missouri. Under Section 40 CFR 130.30, States, Territories or authorized Tribes must submit to the United States Environmental Protection Agency (EPA) a list of impaired or threatened water bodies. - EPA also requires States, Territories, and authorized Tribes to submit a written methodology document describing their approach in considering, and evaluating existing readily available data used to develop their 303(d) list of impaired water bodies. The listing methodology must be submitted to the EPA each year the Section 303(d) list is due. EPA does not approve or disapprove the listing methodology, but considers the methodology during its review of the states 303(d) impaired waters list.

Following the Missouri Clean Water Commission approval, Section 303(d) is submitted to EPA. This fulfills Missouri's biennial submission requirements of an integrated report required under Sections 303(d), 305(b) and 314 of the Clean Water Act. In years when no integrated report is submitted, the department submits a copy of its statewide water quality assessment database to EPA.

### B. U.S. EPA Guidance

In 2001 the Office of General Counsel and the Office of Wetlands, Oceans, and Watersheds developed a recommended framework to assist EPA regions in the preparation of their approval letters for the States' 2002 Section 303(d) list submissions. This was to provide consistency in making approval decisions along with guidance for integrating the development and submission of the 2002 Section 305(b) water quality reports and Section 303(d) list of impaired waters<sup>1</sup>.

The following sections provide an overview of EPA Integrated Report guidance documents from calendar year 2002 through 2015.

The 2002 Integrated Water Quality Monitoring and Assessment Report Guidance was the first document EPA provided to the States, Territories, and authorized Tribes with directions on how to integrate the development and submission of the 2002 305(b) water quality reports and Section 303(d) list of impaired waters.

The guidance recommended that States, Territories and authorized Tribes submit a combined integrated report that would satisfy the Clean Water Act requirements for both Section 305(b) water quality reports and Section 303(d) list. The 2002 Integrated Report was to include:

<sup>1</sup> Additional information can be obtained from EPA's website:  
<http://water.epa.gov/lawsregs/lawguidance/cwa/tmdl/guidance.cfm>.

**Comment [D2]:** Previous comments indicated this section was confusing. This section has been updated to clarify EPA's requirements for the development of a listing methodology.

**Comment [D3]:** Moved this paragraph up to place EPA Guidance discussions in numeric order. The first two sentences were slightly reworded for sentence structure.

Item number 4 was not previously numbered.

Moved reference regarding the methodology is biennially reviewed in order to place it more in sequential order with other information discussed in paragraph.

**Comment [R4]:** This section was updated to add additional background relating to EPA's Integrated Reporting Guidance documents.

Other grammatical changes were also made.

- Delineation of water quality assessment units based on the National Hydrography Dataset (NHD);
- Status of and progress toward achieving comprehensive assessments of all waters;
- Water quality standard attainment status for every assessment units;
- Basis for the water quality standard attainment determinations for every assessment units;
- Additional monitoring that may be needed to determine water quality standard attainment status and, if necessary, to support development of total maximum daily loads (TMDLs) for each pollutant/-assessment units combination;
- Schedules for additional monitoring planned for assessment units;
- Pollutant/-assessment units combinations still requiring TMDLs; and
- TMDL development schedules reflecting the priority ranking of each pollutant/assessment units combination.

The 2002 EPA guidance described the requirements under Section 303(d) of the Clean Water Act where states were required to describe the methodology used to develop their 303(d) list. EPA's guidance recommended the states provide: (1) a description of the methodology used to develop Section 303(d) list; (2) a description of the data and information used to identify impaired and threatened waters; (3) a rationale for not using any readily available data and information; and (4) information on how interstate or international disagreements concerning the list are resolved. Lastly (5), it is recommended that "prior to submission of its Integrated Report, each state should provide the public the opportunity to review and comment on the methodology." In accordance with EPA guidance, the department reviews and updates the Listing Methodology Document (LMD) every two years. The LMD is made available to the public for review and comment at the same time the state's 303(d) impaired waters list is published for public comment. Following the public comment period, the department responds to public comments and provides EPA with a document summarizing all comments received.

In July 2003, EPA issued new guidance entitled "Guidance for 2004 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d) and 305(b) of the Clean Water Act." This guidance gave further recommendations about listing of 303(d) and other waters.

In July 2005, EPA published an amended version entitled "Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act" (see Appendix A for Excerpt).

In October 2006, EPA issued a memorandum entitled "Information Concerning 2008 Clean Water Act Sections 303(d), 305(b) and 314 Integrated Reporting and Listing Decisions." This memorandum serves as EPA's guidance for the 2008 reporting cycle and beyond. This guidance recommended the use of a five-part categorization scheme and that each state provides a comprehensive description of the water quality standards attainment status of all segments within a state (reference Table 1 below). The guidance also defined a "segment" as being used synonymous with the term "assessment unit" used in previous Integrated Report Guidance. Overall, the selected segmentation approach should be consistent with the state's water quality

standards and be capable of providing a spatial scale that is adequate to characterize the water quality standards attainment status for the segment.

It was in the 2006 guidance that EPA recommended all waters of the state be placed in one of five categories described below.

**Table 1. Placement of Waters within the Five Categories in the 2006<sup>2</sup> EPA Assessment, Listing and Reporting Guidance**

<b>Category 1</b>	<p>All <b>designated</b> uses are fully maintained. Data or other information supporting full use attainment for all <b>designated</b> uses must be consistent with the state's Listing Methodology Document (LMD). The department will place a water in Category 1 if the following conditions are met:</p> <ul style="list-style-type: none"><li>• The water has physical and chemical data (at a minimum, water temperature, pH, dissolved oxygen, ammonia, total cobalt, and total copper for streams, and total nitrogen, total phosphorus and secchi depth for lakes) and biological water quality data (at a minimum, <i>E. coli</i> or fecal coliform bacteria) that indicates attainment with water quality standards.</li><li>• The level of mercury in fish fillets or plugs used for human consumption <b>is</b> 0.3 mg/kg or less. Only samples of higher trophic level species (largemouth, smallmouth and spotted bass, sauger, walleye, northern pike, trout, striped bass, white bass, flathead catfish and blue catfish) will be used.</li><li>• The water is not rated as "threatened."</li></ul>
<b>Category 2</b>	<p>One or more designated uses are fully attained but at least one <b>designated</b> use has inadequate data or information to make a use attainment decision consistent with the state's LMD. The department will place a water in Category 2 if at least one of the following conditions are met:</p> <ul style="list-style-type: none"><li>• There <b>is</b> inadequate data for water temperature, pH, dissolved oxygen, ammonia, total cobalt or total copper in streams to assess attainment with water quality standards or inadequate <b>data for</b> total nitrogen, total phosphorus or secchi <b>depth</b> in lakes.</li><li>• There is inadequate <i>E. coli</i> or fecal coliform bacteria data to assess attainment <b>of</b> the whole body contact recreational use.</li><li>• There <b>are</b> insufficient fish fillet tissue, or plug data available for mercury to assess attainment <b>of</b> the fish consumption use.</li></ul> <p>Category 2 waters will be placed in one of two sub-categories.</p> <p>Category 2A: Waters will be placed in this category if available data, using best professional judgement, suggests compliance with <b>numerical</b> water quality criteria of Tables A or B in Missouri's Water</p>

<sup>2</sup> [http://water.epa.gov/lawsregs/lawguidance/cwa/tmdl/upload/2005\\_08\\_11\\_tmdl\\_2006IRG\\_report\\_2006irg-sec5.pdf](http://water.epa.gov/lawsregs/lawguidance/cwa/tmdl/upload/2005_08_11_tmdl_2006IRG_report_2006irg-sec5.pdf)

	<p>Quality Standards (10 CSR 20-7.031) or other quantitative thresholds for determining use attainment.</p> <p>Category 2B: Waters will be placed in this category if the available data, using best professional judgment, suggests noncompliance with numeric water quality criteria of Tables A or B in Missouri’s Water Quality Standards, or other quantitative thresholds for determining use attainment, and <u>these data are</u> insufficient to support a statistical test or to qualify as representative data. Category 2B waters will be given high priority for additional water quality monitoring.</p>
<p><b>Category 3</b></p>	<p>Water quality data are not adequate to assess any of the designated beneficial uses consistent with the LMD. The department will place a water in Category 3 if data are insufficient to support a statistical test or to qualify as representative data to assess any of the designated uses. Category 3 waters will be placed in one of two sub-categories.</p> <p>Category 3A. Waters will be placed in this category if available data, using best professional judgment, suggests compliance with <u>numeric</u> water quality criteria of Tables A or B in Missouri’s Water Quality Standards (10 CSR 20-7.031) or other quantitative thresholds for determining use <u>attainment</u>. Category 3A waters will be tagged for additional water quality monitoring, but will be given lower priority <u>than</u> Category 3B waters.</p> <p>Category 3B. Waters will be placed in this category if the available data, using best professional judgment, suggest noncompliance with <u>numeric</u> water quality criteria of Tables A or B in Missouri’s Water Quality Standards or other quantitative thresholds for determining use attainment. Category 3B waters will be given high priority for additional water quality monitoring.</p>
<p><b>Category 4</b></p>	<p>State <u>water quality standards</u> or other criteria, as per the requirements of <u>Appendix B</u> &amp; C of this document, are not attained, but a Total Maximum Daily Load (TMDL) study is not required. Category 4 waters will be placed in one of three sub-categories.</p> <p>Category 4A. EPA has approved a TMDL study that addresses the impairment. The department will place a water in Category 4A if both the following conditions are met:</p> <ul style="list-style-type: none"> <li>Any portion of the water is rated as being in non-attainment with state <u>water quality standards</u> or other criteria as explained in</li> </ul>

**Comment [D5]:** Since WQ data are not adequate to assess any of the designated uses, EPA suggested that these waters should also be given high priority for additional WQ monitoring

**Comment [R6]:** Previously referenced Table 1

	<p><b>Appendix B &amp; C</b> of this document due to one or more discrete pollutants or discrete properties of the water<sup>3</sup>, and</p> <ul style="list-style-type: none"> <li>EPA has approved a TMDL for all pollutants that are causing non-attainment.</li> </ul> <p>Category 4B. Water pollution controls required by a local, state or federal authority, are expected to correct the impairment in a reasonable period of time. The department will place a water in Category 4B if <b>both</b> of the following conditions are met:</p> <ul style="list-style-type: none"> <li>Any portion of the water is rated as being in non-attainment with state <u>water quality standards</u> or other criteria as explained in <b>Appendix B &amp; C</b> of this document due to one or more discrete pollutants or discrete properties of water<sup>2</sup>, and</li> <li>A water quality based permit that addresses the pollutant(s) causing the designated use, impairment has been issued, and compliance with the permit limits will eliminate the impairment; or other pollution control requirements have been made that are expected to adequately address the pollutant(s) causing the impairment. This may include implemented voluntary watershed control plans as noted in EPA's guidance document.</li> </ul> <p>Category 4C. Any portion of the water is rated as being in non-attainment with state <u>water quality standards</u> or other criteria as explained in <b>Appendix B &amp; C</b> of this document, and a discrete pollutant(s) or other discrete property of the water<sup>2</sup> does not cause the impairment. Discrete pollutants may include specific chemical elements (e.g., lead, zinc), chemical compounds (e.g., ammonia, dieldrin, atrazine) or one of the following quantifiable physical, biological or bacteriological conditions: water temperature, percent of gas saturation, amount of dissolved oxygen, pH, deposited sediment, toxicity or counts of fecal coliform or <i>E. coli</i> bacteria.</p>
<p><b>Category 5</b></p>	<p>At least one discrete pollutant has caused non-attainment with state <u>water quality standards</u> or other criteria as explained in <b>Appendix B &amp; C</b> of this document, and the water does not meet the qualifications for listing as either Categories 4A or 4B. Category 5 waters are those that are candidates for the state's 303(d) List<sup>4</sup>.</p>

Comment [R7]: Previously referenced Table 1

Comment [R8]: Previously referenced Table 1

Comment [R9]: Previously referenced Table 1

Comment [R10]: Previously referenced Table 1

<sup>3</sup> A discrete pollutant or a discrete property of water is defined here as a specific chemical or other attribute of the water (such as temperature, dissolved oxygen or pH) that causes beneficial use impairment and that can be measured quantitatively.

<sup>4</sup> The proposed state 303(d) List is determined by the Missouri Clean Water Commission and the final list is determined by the U.S. Environmental Protection Agency.

	<p>If a designated use is not supported and the segment is impaired or threatened, the fact that a specific pollutant is not known does not provide a basis for excluding a segment from Category 5.</p> <p>Category 5. These segments must be listed as Category 5 unless the state can demonstrate that no discrete pollutant(s) causes or contributes to the impairment. Pollutants causing the impairment will be identified through the 303(d) assessment and listing process before a TMDL study is written. The TMDL should be written within the time frame preferred in EPA guidance for TMDL development, when it fits within the state's TMDL prioritization scheme.</p> <p>Category 5-alt. A water body assigned to 5-alt is an impaired water without a <u>completed</u> TMDL but assigned a low priority for TMDL development because an alternative restoration approach is being pursued. This also provides transparency to the public that a state is pursuing restoration activities in those waters to achieve water quality standards. The addition of this sub-category will facilitate tracking alternative restoration approaches in 303(d) listed waters in priority areas.</p>
<p><b>Threatened Waters</b></p>	<p>When a water is currently attaining all designated uses, but the data shows an inverse (time) trend in quality for one or more discrete water quality pollutants indicating the water will not continue to meet these uses before the next listing cycle. Such water will be considered "threatened." A threatened water will be treated as an impaired water and placed in the appropriate Category (4A, 4B, or 5).</p>

In subsequent years, EPA has provided additional guidance, but only limited new supplemental information has been provided since the 2008 cycle.

In August 2015, the EPA provided draft guidance that would include a Category 5-alternative (5-alt) (reference Table 1 above). Additional information can be found at EPA's website: <http://water.epa.gov/lawsregs/lawsguidance/cwa/tmdl/guidance.cfm>.

**Comment [D11]:** Revised the wording of this definition.

**Comment [D12]:** Moved this so the 2008 and beyond Integrated Report information followed 2006. Any substantial and future Integrated Report guidance information will be added below.

Added 2016 IR information here regarding TMDL prioritization and the additional creation of Category 5-alt.

Moved the "Placement of Waters within the five category in the 2006 EPA Assessment, Listing and Reporting" section into a table so it could be referenced with the text.

## II. The Methodology Document

### A. Procedures and Methods Used to Collect Water Quality Data

#### Department Monitoring

The major purposes of the department's water quality monitoring program are to:

- characterize background or reference water quality conditions;
- better understand daily, flow event and seasonal water quality variations and their underlying processes;
- characterize aquatic biological communities;
- assess time-trends in water quality;
- characterize local and regional effects of point and nonpoint sources pollutants on water quality;
- check for compliance with water quality standards and/or wastewater permit limits;
- support development of strategies, including Total Maximum Daily Loads, to return impaired waters to compliance with Water Quality Standards. All of these objectives are statewide in scope.

#### Coordination with Other Monitoring Efforts in Missouri

To maximize efficiency, the department routinely coordinates its monitoring activities with other agencies to avoid overlap, and to give and receive feedback on monitoring design. Data from other sources are used for meeting the same objectives as department-sponsored monitoring. The data must fit the criteria described in the data quality considerations section of this document. The agencies most often involved are the U.S. Geological Survey, the U.S. Army Corps of Engineers, EPA, the Missouri Department of Conservation (MDC), and the Missouri Department of Health and Senior Services. The Department of Natural Resources also tracks the monitoring efforts of the National Park Service; the U.S. Forest Service; several of the state's larger cities; the states of Oklahoma, Arkansas, Kansas, Iowa, and Illinois; and graduate level research conducted at universities within Missouri. For those wastewater discharges where the department has required instream water quality monitoring, the department may also use monitoring data acquired by wastewater dischargers as a condition of discharge permits issued by the department. In 1995, the department also began using data collected by volunteers that have passed Volunteer Water Quality Monitoring Program Quality Assurance/Quality Control tests.

#### Existing Monitoring Networks and Programs

The following is a list and a brief description of the kinds of water quality monitoring activities presently occurring in Missouri.

## 1. Fixed Station Network

- a) Objective: To better characterize background or reference water quality conditions, to better understand daily, flow events, and seasonal water quality variations and their underlying processes, to assess trends and to check for compliance with water quality standards.
- b) Design Methodology: Sites are chosen based on one of the following criteria:
  - Site is believed to have water quality representative of many neighboring streams of similar size due to similarity in watershed geology, hydrology and land use, and the absence of any impact from a significant point or discrete nonpoint water pollution source.
  - Site is downstream of a significant point source or discrete nonpoint source area.
- c) Number of Sites, Sampling Methods, Sampling Frequency, and Parameters:
  - MDNR/U.S. Geological Survey cooperative network: approximately 70 sites statewide, horizontally and vertically integrated grab samples, four to twelve times per year. Samples are analyzed for major ions (e.g. calcium, magnesium, sulfate, and chloride), nutrients (e.g. phosphorus and nitrogen), temperature, pH, dissolved oxygen, specific conductance, bacteria (e.g. E. coli and fecal coliform) and flow on all visits, two to four times annually for suspended solids and heavy metals, and for pesticides six times annually at four sites.
  - MDNR/University of Missouri-Columbia's lake monitoring network. This program has monitored about 249 lakes since 1989. About 75 lakes are monitored each year. Each lake is usually sampled four times during the summer and about 12 are monitored spring through fall for nutrients, chlorophyll, turbidity and suspended solids.
  - Department routine monitoring of finished public drinking water supplies for bacteria and trace contaminants.
  - Routine bacterial monitoring for E. coli of swimming beaches at Missouri's state parks during the recreational season by the department's Missouri State Parks.
  - Monitoring of sediment quality by the department at approximately 10-12 discretionary sites annually. Sites are monitored for several heavy metals (e.g. arsenic, cadmium, copper, lead, mercury, nickel, zinc, etc.) and/or organic contaminants (e.g. polycyclic aromatic hydrocarbons, etc.).

## 2. Special Water Quality Studies

- a) Objective: Special water quality studies are used to characterize water quality effects from a specific pollutant source area.

b) Design Methodology: These studies are designed to verify and measure the contaminants of concern based on previous water quality studies, effluent sampling and/or Missouri State Operating Permit applications. These studies employ multiple sampling stations downstream and upstream (if appropriate). If contaminants of concern have significant seasonal or daily variation, the sampling design must account for such variation.

Comment [D13]: Streamlined sentence wording

c) Number of Sites, Sampling Methods, Sampling Frequency and Parameters: The department conducts or contracts up to 10 to 15 special studies annually, as funding allows. Each study has multiple sampling sites. The number of sites, sampling frequency and parameters all vary greatly depending on the study. Intensive studies would also require multiple samples per site over a relatively short time frame.

### 3. Toxics Monitoring Program

The fixed station network and many of the department's intensive studies monitor for acute and chronic toxic chemicals<sup>5</sup>. In addition, major municipal and industrial dischargers must monitor for acute and chronic toxicity in their effluents as a condition of their Missouri State Operating Permit.

### 4. Biological Monitoring Program

a) Objectives: The objectives of the Biological Monitoring programs are to develop numeric criteria describing "reference" aquatic macroinvertebrate and fish communities in Missouri's streams, to implement these criteria within state water quality standards and to maintain a statewide fish and aquatic macroinvertebrate monitoring program.

b) Design Methodology: Development of biocriteria for fish and aquatic macroinvertebrates<sup>6</sup> involves identification of reference streams in each of Missouri's aquatic ecoregions and 17 ecological drainage units, respectively. It also includes intensive sampling of invertebrate and fish communities to quantify temporal and spatial variation in reference streams within ecoregions and variation among ecoregions, and the sampling of chemically and physically impaired streams to test sensitivity of various community metrics to differences in stream quality.

c) Number of Sites, Sampling Methods, Sampling Frequency and Parameters: The department has conducted biological sampling of aquatic macroinvertebrates for many years. Since 1991, the department's aquatic macroinvertebrate monitoring program has consisted of standardized monitoring of approximately 45 to 55 sites twice annually. In addition, the MDC presently has a statewide fish and aquatic macroinvertebrate monitoring program, the Resource Assessment and Monitoring (RAM) Program, designed monitor and assess the health of Missouri's stream resources on a rotating basis. This program samples a minimum of 450 random and 30 reference sites every five years.

<sup>5</sup> As defined in 10 CSR 20-7.031(1)

<sup>6</sup> For additional information visit: <http://dnr.mo.gov/env/esp/wqm/biologicalassessments.htm>

## 5. Fish Tissue Monitoring Program

- a) Objective: Fish tissue monitoring addresses two objectives: (1) the assessment of ecological health or the health of aquatic biota (usually accomplished by monitoring whole fish samples); and (2) the assessment of human health risk based on the level of contamination of fish tissue plugs, or fillets.
- b) Design Methodology: Fish tissue monitoring sites are chosen based on one of the following criteria:
  - Site is believed to have water and sediment quality representative of many neighboring streams or lakes of similar size due to similarity in geology, hydrology and land use, and the absence of any known impact from a significant point source or discrete nonpoint water pollution source.
  - Site is downstream of a significant point source or discrete nonpoint source area.
  - Site has shown fish tissue contamination in the past.
- c) Number of Sites, Sampling Methods, Sampling Frequency and Parameters:

The department plans to maintain a fish tissue monitoring program to collect whole fish composite samples<sup>7</sup> at approximately 13 fixed sites. In previous years, this was a cooperative effort between EPA and the department through EPA's Regional Ambient Fish Tissue (RAFT) Monitoring Program. Each site will be sampled once every two years. The preferred species for these sites are either Common Carp (*Cyprinus carpio*) or one of the Redhorse (a.k.a. sucker) species (*Moxostoma* sp.).

The department, EPA, and MDC also sample 40 to 50 discretionary sites annually for two fish fillet composite samples or fish tissue plug samples (mercury only) from fish of similar size and species. One sample is of a top carnivore such as Largemouth Bass (*Micropterus salmoides*), Smallmouth Bass (*Micropterus dolomieu*), Walleye (*Zander vitreus*), or Sauger (*Zander lucioperca*). The other sample is for a species of a lower trophic level such as catfish, Common Carp or sucker species (*Catostomidae*). This program occasionally samples fish eggs for certain fish species at selected locations. Both of these monitoring programs analyze for several chlorinated hydrocarbon insecticides, PCBs, lead, cadmium, mercury, and fat content.

## 6. Volunteer Monitoring Program

Two major volunteer monitoring programs generate water quality data in Missouri. The data generated from these programs are used for statewide 305(b) reporting on general water quality health, used as a screening level tool to determine where additional monitoring is needed, or used to supplement other water quality data for watershed planning purposes.

<sup>7</sup> A composite sample is one in which several individual fish are combined to produce one sample.

- Lakes of Missouri Volunteer Program<sup>8</sup>. This cooperative program consists of persons from the department, the University of Missouri-Columbia, and volunteers who monitor approximately 137 sites on 66 lakes, including Lake Taneycomo, Table Rock Lake and several lakes in the Kansas City area. Lake volunteers are trained to collect samples for total phosphorus, total nitrogen, chlorophyll and inorganic suspended sediments. Data from this program is used by the university as part of a long-term study on the limnology of mid-western reservoirs.
- Volunteer Water Quality Monitoring Program. The Volunteer Water Quality Monitoring Program<sup>9</sup> is an activity of the Missouri Stream Team Program, which is a cooperative project sponsored by the department, the Missouri Department of Conservation, and the Conservation Federation of Missouri. The program involves volunteers who monitor water quality of streams throughout Missouri. There are currently over 5,000 Stream Teams and more than 3,600 trained water quality monitors. Approximately 80,000 citizens are served each year through the program. Since the beginning of the Stream Team program, 494,232 volunteers have donated about 2 million hours valued at more than \$38 million to the State of Missouri.

After the Introductory class, many attend at least one more class of higher level training: Levels 1, 2, 3 and 4. Each level of training is a prerequisite for the next higher level, as is appropriate data submission. Data generated by Levels 2, 3, and 4 and the new Cooperative Site Investigation (CSI) Program volunteers represent increasingly higher quality assurance. For CSI projects, the volunteers have completed a quality assurance/quality control workshop, completed field evaluation, and/or have been trained to collect samples following department protocols. Upon completing Introductory and Level 1 and 2 training, volunteers will have received the basic level training to conduct visual stream surveys, stream discharge measurements, biological monitoring, and collect physical and chemical measurements for pH, conductivity, dissolved oxygen, nitrate, phosphate, and turbidity.

Of those completing an Introductory course, about 35 percent proceed to Levels 1 and 2. To date, 104 volunteers have reached Level 3 and six volunteers have reached Level 4. The CSI Program uses trained volunteers to collect samples and transport them to laboratories approved by the department. Volunteers and department staff work together to develop a monitoring plan. Currently there are 39 volunteers qualified to work in the CSI Program. All Level 2, 3, and 4 volunteers, as well as all CSI trained volunteers, are required to attend a validation session every 3 years to ensure equipment, reagents and methods meet program standards. To date 106 individuals have attended a validation session at least once.

<sup>8</sup> For additional program information visit: <http://www.lmvp.org/>

<sup>9</sup> For additional program information visit: <http://dnr.mo.gov/env/wpp/VWQM.htm>

## Identification of All Existing and Readily Available Water Quality Data Sources

### Data Solicitation Request

Comment [R14]: Added an overview of the Data Solicitation Process

In calendar year 2014, the department sent out a request for all available water quality data (chemical and biological). The data solicitation requested water quality data for approximately a two year timeframe prior to the current listing year. The data solicitation request was sent to multiple agencies, neighboring states, and organizations. In addition, and as part of the data solicitation process, the department queries available water quality data from national databases such as EPA's Storage and Retrieval (STORET)/Water Quality Exchange (WQX) data warehouse<sup>10</sup>, and the USGS Water Quality Portal<sup>11</sup>.

The data must be spatially and temporally representative of the actual annual ambient conditions of the water body. Sample locations should be characteristic and representative of the main water mass or distinct hydrologic areas. With the exception of the data collected for those designated uses that require seasonally based data (e.g., whole body contact recreation, biological community data, and critical season dissolved oxygen), data should be distributed over at least three seasons, over two years, and should not be biased toward specific conditions (such as runoff, season, or hydrologic conditions).

Data meeting the following criteria will be accepted.

- Samples must be collected and analyzed under a Quality Assurance/Quality Control (QA/QC) protocol that follows the EPA requirements for quality assurance project plans.
- Samples must be analyzed following protocols that are consistent with the EPA or Standard Method procedures.
- All data submitted must be accompanied by a copy of the organization's QA/QC protocol and standard operating procedures.
- All data must be reported in standard units as recommended in the relevant approved methods.
- All data must be accompanied by precise sample location(s), preferably in either decimal degrees or Universal Transverse Mercator (UTM).
- All data must be received in a Microsoft Excel or compatible format.
- All data must have been collected within the requested period of record.

All readily available and acceptable data are uploaded into the department's Water Quality Assessment Database<sup>12</sup>, where the data undergoes quality control checks prior to 303(d) or 305(b) assessment processes.

<sup>10</sup> [http://www.epa.gov/storet/dw\\_home.html](http://www.epa.gov/storet/dw_home.html)

<sup>11</sup> <http://www.waterqualitydata.us/>

<sup>12</sup> [http://dnr.mo.gov/mocwis\\_public/wqa/waterbodySearch.do](http://dnr.mo.gov/mocwis_public/wqa/waterbodySearch.do)

### Laboratory Analytical Support

#### Laboratories used:

- Department/U.S. Geological Survey Cooperative Fixed Station Network: U.S. Geological Survey Lab, Denver, Colorado
- Intensive Surveys: Varies, many are done by the department's Environmental Services Program
- Toxicity Testing of Effluents: Many commercial laboratories
- Biological Criteria for Aquatic Macroinvertebrates: department's Environmental Services Program and University of Missouri-Columbia
- Fish Tissue: EPA Region VII Laboratory, Kansas City, Kansas, and miscellaneous contract laboratories (Missouri Department of Conservation or U.S. Geological Survey's Columbia Environmental Research Center)
- Missouri State Operating Permit: Self-monitoring or commercial laboratories
- Department's Public Drinking Water Monitoring: department's Environmental Services Program and commercial laboratories<sup>13</sup>
- Other water quality studies: Many commercial laboratories

### **B. Sources of Water Quality Data**

The following data sources are used by the department to aid in the compilation of the state's integrated report (previously the 305(b) report). Where quality assurance programs are deemed acceptable, additional sources would also be used to develop the state's Section 303(d) list. These sources presently include, but are not limited to:

1. Fixed station water quality and sediment data collected and analyzed by the department's Environmental Services Program personnel.
2. Fixed station water quality data collected by the U.S. Geological Survey under contractual agreements with the department.
3. Fixed station water quality data collected by the U.S. Geological Survey under contractual agreements to agencies or organizations other than the department.
4. Fixed station water quality, sediment quality, and aquatic biological information collected by the U.S. Geological Survey under their National Stream Quality Accounting Network and the National Water Quality Assessment Monitoring Programs.
5. Fixed station raw water quality data collected by the Kansas City Water Services Department, the St. Louis City Water Company, the Missouri American Water Company (formerly St. Louis County Water Company), Springfield City Utilities, and Springfield's Department of Public Works.
6. Fixed station water quality data collected by the U.S. Army Corps of Engineers. The

<sup>13</sup> For additional information visit: <http://dnr.mo.gov/env/wpp/labs/>

- Kansas City, St. Louis, and Little Rock Corps Districts have monitoring programs for Corps-operated reservoirs in Missouri.
7. Fixed station water quality data collected by the Arkansas Department of Environmental Quality, the Kansas Department of Health and Environment, the Iowa Department of Natural Resources, and the Illinois Environmental Protection Agency.
  8. Fixed station water quality monitoring by corporations.
  9. Annual fish tissue monitoring programs by [EPA/Department RAFT Monitoring Program](#) and [MDC](#).
  10. Special water quality surveys conducted by the [department](#). Most of these surveys are focused on the water quality impacts of specific point source wastewater discharges. Some surveys are of well-delimited nonpoint sources such as abandoned mined lands. These surveys often include physical habitat evaluation and monitoring of aquatic [macroinvertebrates](#) as well as water chemistry monitoring.
  11. Special water quality surveys conducted by U.S. Geological Survey, including but not limited to:
    - a) Geology, hydrology and water quality of various hazardous waste sites,
    - b) Geology, hydrology and water quality of various abandoned mining areas,
    - c) Hydrology and water quality of urban nonpoint source runoff in [metropolitan areas of Missouri \(e.g. St. Louis, Kansas City, and Springfield\)](#), and
    - d) Bacterial and nutrient contamination of streams in southern Missouri.
  12. Special water quality studies by other agencies such as [MDC](#), the U.S. Public Health Service, and the Missouri Department of Health and Senior Services.
  13. Monitoring of fish occurrence and distribution by [MDC](#).
  14. Fish Kill and Water Pollution Investigations Reports published by [MDC](#).
  15. Selected graduate research projects pertaining to water quality and/or aquatic biology.
  16. Water quality, sediment, and aquatic biological data collected by the [department](#), [EPA](#) or their contractors at hazardous waste sites in Missouri.
  17. Self-monitoring of receiving streams by cities, sewer districts and industries, or contractors on their behalf, for those discharges that require this kind of monitoring. This monitoring includes chemical and sometimes toxicity monitoring of some of the larger wastewater discharges, particularly those that discharge to smaller streams and have the greatest potential to affect instream water quality.
  18. Compliance monitoring of receiving waters by the [department](#) and EPA. This can include chemical and toxicity monitoring.
  19. Bacterial monitoring of streams and lakes by county health departments, community lake associations, and other organizations using acceptable analytical methods.
  20. Other monitoring activities done under a quality assurance project plan approved by

the department.

21. Fixed station water quality and aquatic macroinvertebrate monitoring by volunteers who have successfully completed the Volunteer Water Quality Monitoring Program Level 2 workshop. Data collected by volunteers who have successfully completed a training Level 2 workshop is considered to be Data Code One. Data generated from Volunteer Training Levels 2, 3 and 4 are considered "screening" level data and can be useful in providing an indication of a water quality problem. For this reason, the data are eligible for use in distinguishing between waters in Categories 2A and 2B or Categories 3A and 3B. Most of this data are not used to place waters in main Categories (1, 2, 3, 4, and 5) because analytical procedures do not use EPA or Standard Methods or other department approved methods. Data from volunteers who have not yet completed a Level 2 training workshop do not have sufficient quality assurance to be used for assessment. Data generated by volunteers while participating in the department's Cooperative Site Investigation Program (Section II C1) or other volunteer data that otherwise meets the quality assurance outlined in Section II C2 may be used in Section 303(d) assessment.

The following data sources (22-23) cannot be used to rate a water as impaired (Categories 4A, 4B, 4C or 5); however, these data sources may be used to direct additional monitoring that would allow a water quality assessment for Section 303(d) listing.

22. Fish Management Basin Plans published by MDC.
23. Fish Consumption Advisories published annually by the Missouri Department of Health and Senior Services. Note: the department may use data from data source listed as Number 9 above, to list individual waters as impaired due to contaminated fish tissue.

As previously stated, the department will review all data of acceptable quality that are submitted to the department prior to the first public notice of the draft 303(d) list. However, the department will reserves the right to review and use data of acceptable quality submitted after this date if the data results in a change to the assessment outcome of the water.

### C. Data Quality Considerations

- DNR Quality Assurance/Quality Control Program

The department and EPA Region VII have completed a Quality Management Plan. All environmental data generated directly by the department, or through contracts funded by the department, or EPA require a Quality Assurance Project Plan<sup>14</sup>. The agency or organization responsible for collecting and/or analyzing environmental data must write and adhere to a Quality Assurance Project Plan approved through the department's

<sup>14</sup> For additional information visit: <http://www.epa.gov/quality/gapps.html>

Quality Management Plan. Any environmental data generated via a monitoring plan with a department approved Quality Assurance Project Plan are considered suitable for use in water quality assessment and the 303(d) listing. This includes data generated by volunteers participating in the department's CSI Program. Under this program, the department's Environmental Services Program will audit selected non-profit (governmental and university) laboratories. Laboratories that pass this audit will be approved for the CSI Program. Individual volunteers who collect field samples and deliver them to an approved laboratory must first successfully complete department training on how to properly collect and handle environmental samples. The types of information that will allow the department to make a judgment on the acceptability of a quality assurance program are: (1) a description of the training, and work experience of the persons involved in the program, (2) a description of the field meters and maintenance and calibration procedures, (3) a description of sample collection and handling procedures, and (4) a description of all analytical methods used in the laboratory for analysis.

- Other Quality Assurance/Quality Control Programs

Data generated in the absence of a department-approved Quality Assurance Project Plan may be used to assess a water body if the department determines that the data are adequate after reviewing and accepting the quality assurance procedures plan used by the data generator. This review would include: (1) names of all persons involved in the monitoring program, their duties, and a description of their training and work related experience, (2) all written procedures, Standard Operating Procedures, or Quality Assurance Project Plans pertaining to this monitoring effort, (3) a description of all field methods used, brand names and model numbers of any equipment, and a description of calibration and maintenance procedures, and (4) a description of laboratory analytical methods. This review may also include an audit by the department's Environmental Services Program.

- Other Data Quality Considerations

3.1 Data Age. For assessing present conditions, more recent data are preferable; however, older data may be used to assess present conditions if the data remains representative of present conditions.

If the department uses data older than seven years to make a Section 303(d) list decision a written justification for the use of such data will be provided.

A second consideration is the age of the data relative to significant events that may have an effect on water quality. Data collected prior to the initiation, closure, or significant change in a wastewater discharge, or prior to a large spill event or the reclamation of a mining or hazardous waste site, for example, may not be representative of present conditions. Such data would not be used to assess present conditions even if it was less

than seven years old. Such “pre-event” data can be used to determine changes in water quality before and after the event or to show water quality trends.

3.2 Data Type, Amount and Information Content. EPA recommends establishing a series of data codes, and rating data quality by the kind and amount of data present at a particular location (EPA 1997<sup>15</sup>). The codes are single-digit numbers from one to four, indicating the relative degree of assurance the user has in the value of a particular environmental data set. Data Code One indicates the least assurance or the least number of samples or analytes and Data Code Four the greatest. Based on EPA’s guidance, the department uses the following rules to assign code numbers to data.

- Data Code<sup>16</sup> One: All data not meeting the requirements of the other data codes.
- Data Code Two: Chemical data collected quarterly to bimonthly for at least three years, or intensive studies that monitor several nearby sites repeatedly over short periods of time, or at least three composite or plug fish tissue samples per water body, or at least five bacterial samples collected during the recreational season of one calendar year.
- Data Code Three: Chemical data collected at least monthly for more than three years on a variety of water quality constituents including heavy metals and pesticides; or a minimum of one quantitative biological monitoring study of at least one aquatic assemblage (fish, macroinvertebrates, or algae) at multiple sites, or multiple samples at a single site when data from that site is supported by biological monitoring at an appropriate control site.
- Data Code Four: Chemical data collected at least monthly for more than three years that provides data on a variety of water quality constituents including heavy metals and pesticides, and including chemical sampling of sediments and fish tissue; or a minimum of one quantitative biological monitoring study of at least two aquatic assemblages (fish, macroinvertebrates, or algae) at multiple sites.

In Missouri, the primary purpose of Data Code One data is to provide a rapid and inexpensive method of screening large numbers of waters for obvious water quality problems and to determine where more intensive monitoring is needed. In the preparation of the state’s Integrated Report, data from all four data quality levels are used. Most of the data is of Data Code One quality, and without Data Code One data, the department would not be able to assess a majority of the state’s waters.

<sup>15</sup> *Guidelines for the Preparation of the Comprehensive State Water Quality Assessments (305b) and Electronic Updates*, 1997. (<http://water.epa.gov/type/watersheds/monitoring/repguid.cfm>)

<sup>16</sup> Data Code One is equivalent to data water quality assurance Level One in 10 CSR 20-7.050 General Methodology for Development of Impaired Waters List, subsection (2)(C), Data Code Two is equivalent to Level 2, etc.

In general, when selecting water bodies for the Missouri 303(d) List, only Data Code Two or higher are used, unless the problem can be accurately characterized by Data Code One data.<sup>17</sup> The reason is that Data Code Two data provides a higher level of assurance that a Water Quality Standard is not actually being attained and that a TMDL study is necessary. All water bodies placed in Categories 2B or 3B receive high priority for additional monitoring so that data quality is upgraded to at least Data Code Two. Category 2B and 3B waters will be given higher priority than Categories 2A and 3A.

#### D. How Water Quality Data is Evaluated to Determine Whether or Not Waters are Impaired for 303(d) Listing Purposes

- Physical, Chemical, Biological and Toxicity Data

During each reporting cycle, the department and stakeholders review and revise the guidelines for determining water quality impairment. The guidelines shown in Appendix B & C provides the general rules of data use and assessment and Appendix D provides details about the specific analytical procedure used. In addition, if trend analysis indicates that presently unimpaired waters will become impaired prior to the next listing cycle, these “threatened waters” will be judged as impaired. Where antidegradation provisions in Missouri’s Water Quality Standards apply, those provisions shall be upheld. The numerical criteria included in Appendix B have been adopted into the state water quality standards, 10 CSR 20-7.031, and are used, as described in Appendix B to make use attainment decisions.

- Weight of Evidence Approach

When evaluating narrative criteria described in the state water quality standards, 10 CSR 20-7.031, the department will use a weight of evidence analysis for assessing numerical translators that have not been adopted into state water quality standards (see Appendix C). Under the weight of evidence approach, all available information is examined and the greatest weight is given to data providing the “best supporting evidence” for an attainment decision. Determination of “best supporting evidence” will be made using best professional judgment, considering factors such as data quality, and site-specific environmental conditions. This weight of evidence analysis will include the use of other types of environmental data when it is available or collection of additional data to make the most informed use attainment decision. Examples of other relevant environmental data might include biological data on fish [Fish Index of Biotic Integrity (fBI)] or aquatic macroinvertebrate [Macroinvertebrate Stream Condition Index (MSCI)] scores, fish tissue, or toxicity testing of water or sediments.

<sup>17</sup> When a listing, amendment or delisting of a 303(d) water is made with only Data Code One data, a document will be prepared that includes a display of all data and a presentation of all statistical tests or other evaluative techniques that documents the scientific defensibility of the data. This requirement applies to all Data Code One data identified in Appendix B of this document.

Comment [R15]: Previously Tables 1.1 and 1.2

Comment [R16]: Previously Table 2.1

Comment [R17]: Previously Table 1

Comment [R18]: Previously Table 1.1

Comment [D19]: The previous Appendix E wording was moved to this section and blended with previous wording. Repetitive wording was removed.

Biological data will be given greater weight in a weight of evidence analysis for making attainment decisions for aquatic life use and subsequent Section 303(d) listings. Whether or not numeric translators of biological criteria are met is a strong indicator for the attainment of aquatic life use. Moreover, the department retains a high degree of confidence in an attainment decision based on biological data that is representative of water quality condition.

When the weight of evidence analysis suggests, but does not provide strong scientifically valid evidence of impairment, the department will place the water body in question in Categories 2B or 3B. The department will produce a document showing all relevant data and the rationale for the attainment decision. All such documents will be available to the public at the time of the first public notice of the proposed 303(d) list. A final recommendation on the listing of a water body based on narrative criteria will only be made after full consideration of all comments on the proposed list.

- **Biological Data**

Methods for assessing biological data typically receive considerable attention during the public comment period of development of the Listing Methodology Documents. Currently, a defined set of biocriteria are used to evaluate biological data for assessing compliance with water quality standards. These biological criteria contain numeric thresholds, that when exceeded relative to prescribed assessment methods, serve as a basis for identifying candidate waters for Section 303(d) listing. Biocriteria are based on three types of biological data, including: (1) aquatic macroinvertebrate community data; (2) fish community data; and, (3) a catch-all class referred to as “other biological data.”

In general, for interpretation of macroinvertebrate data, where habitat assessment scores indicate habitat is less than 75 percent of reference or appropriate control stream scores, and in the absence of other data indicating impairment by a discrete pollutant, a water body judged to be impaired will be placed in Category 4C. When interpreting fish community data, a provisional multi-metric habitat index called the QCPHI index is used to identify stream habitat in poor condition. The QCPHI index separates adequate habitat from poor habitat using a 0.39 threshold value; whereby, QCPHI scores < 0.39 indicate stream habitat is of poor quality, and scores greater than 0.39 indicate available stream habitat is adequate. In the absence of other data indicating impairment by a discrete pollutant, impaired fish communities with poor habitat will be placed in Category 4C. Additional information about QCPHI is provided in the Considerations for the Influence of Habitat Quality and Sample Representativeness section.

The sections below describe the methods used to evaluate the three types of biological data (macroinvertebrate community, fish community, and other biological data), along with background information on the development and scoring of biological criteria, procedures for assessing biological data, methods used to ensure sample

**Comment [D20]:** Blend text with the previous Appendix E. Any repetitive wording removed.

representativeness, and additional information used to aid in assessing biological data such as the weight of evidence approach.

#### *Aquatic Macroinvertebrate Community Data*

The department conducts aquatic biological assessments to determine macroinvertebrate community health as a function of water quality and habitat. The health of a macroinvertebrate community is directly related to water quality and habitat. Almost all macroinvertebrate evaluation consists of comparing the health of the community of the “target” to healthy macroinvertebrate communities from reference streams of the same general size and usually in the same Ecological Drainage Unit (EDU).

The department’s approach to monitoring and evaluating aquatic macroinvertebrates is largely based on *Biological Criteria for Wadeable/Perennial Streams of Missouri* (MDNR 2002). This document provides numerical biological criteria (biocriteria) relevant to the protection of aquatic life use for wadeable streams in the state. Biocriteria were developed using wadeable reference streams that occur in specific EDUs as mapped by the Missouri Resource Assessment Partnership. For macroinvertebrates, the numerical biocriterion translator is expressed as a multiple metric index referred to as the MSCI. The MSCI includes four metrics: Taxa Richness (TR); Ephemeroptera, Plecoptera, and Trichoptera Taxa (EPTT); Biotic Index (BI); and the Shannon Diversity Index (SDI). These metrics are considered indicators of stream health, and change predictably in response to the environmental condition of a stream.

Metric values are determined directly from macroinvertebrate sampling. To calculate the MSCI, each metric is normalized to unitless values of 5, 3, or 1, which are then added together for a total possible score of 20. MSCI scores are divided into three levels of stream condition:

- Fully Biologically Supporting (16-20),
- Partially Biologically Supporting (10-14), and
- Non-Biologically Supporting (4-8).

Partially and Non-Biologically Supporting streams may be considered impaired and are candidates for Section 303(d) listing.

Unitless metric values (5, 3, or 1) were developed from the lower quartile of the distribution of each metric as calculated from reference streams for each EDU. The lower quartile (25<sup>th</sup> percentile) of each metric equates to the minimum value still representative of unimpaired conditions. In operational assessments, metric values below the lower quartile of reference conditions are typically judged as impaired (United States Environmental Protection Agency 1996, Ohio Environmental Protection Agency 1990, Barbour *et al.* 1996). Moreover, using the 25<sup>th</sup> percentile of reference conditions for each metric as a

standard for impairment allows natural variability to be filtered out. For metrics with values that decrease with increasing impairment (TR, EPTT, SDI), any value above the lower quartile of the reference distribution receives a score of five. For the BI, whose value increases with increasing impairment, any value below the upper quartile (75<sup>th</sup> percentile) of the reference distribution receives a score of five. The remainder of each metric's potential quartile range below the lower quartile is bisected, and scored either a three or a one. If the metric value is less than or equal to the quartile value and greater than the bisection value it is scored a three. If the metric value is less than or equal to the bisection value it is scored a one.

MSCI scores meeting data quality considerations may be assessed for the protection of aquatic life using the following procedures.

**Determining Full Attainment of Aquatic Life Use:**

- For seven or fewer samples, 75% of the MSCI scores must be 16 or greater. Fauna achieving these scores are considered to be very similar to biocriteria reference streams.
- For eight or more samples, results must be statistically similar to representative reference or control streams.

**Determining Non-Attainment of Aquatic Life Use:**

- For seven or fewer samples, 75% of the MSCI scores must be 14 or lower. Fauna achieving these scores are considered to be substantially different from biocriteria reference streams.
- For eight or more samples, results must be statistically dissimilar to representative reference or control streams.

**Data will be judged inconclusive when outcomes do not meet requirements for decisions of full or non-attainment.**

As noted, when eight or more samples are available, results must be statistically similar or dissimilar to reference or control conditions in order to make an attainment decision. To accomplish this, a binomial probability with an appropriate level of significance ( $\alpha$ =alpha), is calculated based on the null hypothesis that the test stream would have a similar percentage of MSCI scores that are 16 or greater as reference streams. The significance level is set at  $\alpha=0.1$ , meaning if the p-value of the hypothesis test is less than  $\alpha$ , the hypothesis is considered statistically significant. The significance level of  $\alpha$  is in fact the probability of making a wrong decision and committing a Type I error (rejecting a true null hypothesis). When the Type I error rate is less than  $\alpha=0.1$ , the null hypothesis is rejected. Inversely, when the Type I error rate is greater than  $\alpha=0.1$ , the null hypothesis is accepted. For comparing samples from a test stream to samples collected from reference streams in the same EDU, the percentage of samples from reference streams scoring 16 or greater is used to determine the probability of "success" and "failure" in the

binomial probability equation. For example, if 84% of the reference stream MSCI scores in a particular EDU are 16 or greater, then 0.84 would be used as the probability of success and 0.16 would be used as the probability of failure. Note that [Appendix I](#) states to “rate a stream as impaired if biological criteria reference stream frequency of fully biologically supporting scores is greater than five percent more than the test stream,” thus, a value of 0.79 (0.84 - 0.05) would actually be used as the probability of success in the binomial distribution equation.

Comment [R21]: Previously Table B-1

**Binomial Probability Example:**

Reference streams from the Ozark/Gasconade EDU classified as riffle/pool stream types with warm water temperature regimes produce fully biologically supporting streams 85.7% of the time. In the test stream of interest, six out of ten samples resulted in MSCI scores of 16 or more. Calculate the Type I error rate for the probability of getting six or fewer fully biologically supporting scores in ten samples.

The binomial probability formula may be summarized as:

$$p^n + (n!/ X!(n-X)! * p^n q^{n-x}) = 1$$

Where,

Sample Size (n) = 10

Number of Successes (X) = 6

Probability of Success (p) = 0.857 - 0.05 = 0.807

Probability of Failure (q) = 0.193

Binomial Distribution Coefficients = n!/ X!(n-X)!

The equation may then be written as:

$$\begin{aligned} &= 1 - ((0.807^{10}) + ((10*(0.807^9)*(0.193))) + ((45*(0.807^8)*(0.193^2)) + \\ &((120*(0.807^7) * (0.193^3))) \\ &= 0.109 \end{aligned}$$

Since 0.109 is greater than the test significance level (minimum allowable Type I error rate) of  $\alpha=0.1$ , we accept the null hypothesis that the test stream has the same percent of fully biologically supporting scores as the same type of reference streams from the Ozark/Gasconade EDU. Thus, this test stream would be judged as unimpaired.

If under the same scenario, there were only 5 samples from the test stream with MSCI scores of 16 or greater, the Type I error rate would change to 0.028, and since this value is less than the significance level of  $\alpha=0.1$ , the stream would be judged as impaired.

Within each EDU, MSCI scores are categorized by sampling regime (Glide/Pool vs. Riffle/Pool) and temperature regime (warm water vs. cold water). The percentage of fully biologically supporting scores for the Mississippi River Alluvial Basin/Black/Cache EDU is not available due to the lack of reference sites in this region. Percentages of fully biologically supporting samples per EDU is not included here, but can be made available upon request. The percentage of reference streams per EDU that are fully biologically supporting may change periodically as additional macroinvertebrate samples are collected and processed from reference samples within an EDU.

#### ***Sample Representativeness***

The departments field and laboratory methods used to collect and process macroinvertebrate samples are contained in the document *Semi-Quantitative Macroinvertebrate Stream Bioassessment* (MDNR 2012a). Macroinvertebrates are identified to levels following standard operating procedures contained in *Taxonomic Levels for Macroinvertebrate Identifications* (MDNR 2012b). Macroinvertebrate monitoring is accompanied by physical habitat evaluations as described in the document *Stream Habitat Assessment* (MDNR 2010). For the assessment of macroinvertebrate samples, available information must meet data code levels three and four as described in Section II.C of this LMD. Data coded as levels three and four represent environmental data providing the greatest degree of assurance. Thus, at a minimum, macroinvertebrate assessments include multiple samples from a single site, or samples from multiple sites within a single reach.

It is important to avoid situations where poor or inadequate habitat prohibits macroinvertebrate communities from being assessed as fully biologically supporting. Therefore, when assessing macroinvertebrate samples, the quality of available habitat must be similar to that of reference streams within the appropriate EDU. The department's policy for addressing this concern has been to exclude MSCI scores from an assessment when accompanying habitat scores are less than 75 percent of the mean habitat scores from reference streams of the appropriate EDU. The following procedures outline the department's method for assessing macroinvertebrate communities from sites with poor or inadequate habitat.

#### ***Assessing Macroinvertebrate Communities from Poor/Inadequate Habitat:***

- If less than half the macroinvertebrate samples in an assessed stream segment have habitat scores less than 75 percent of the mean score for reference streams in that EDU, any sample that scores less than 16 and has a habitat score less than 75 percent of the mean reference stream score for that EDU, is excluded from the assessment process.
- If at least half the macroinvertebrate samples in an assessed stream segment have habitat scores less than 75 percent of the mean score for reference streams in that EDU and the assessment results in a judgment that the

macroinvertebrate community is impaired, the assessed segment will be placed in Category 4C impairment due to poor aquatic habitat.

- If one portion of the assessment reach contains two or more samples with habitat scores less than 75 percent of reference streams from that EDU while the remaining portion does not, the portion of the stream with poor habitat scores could be separately assessed as a category 4C stream permitting low MSCI scores.

Macroinvertebrate sampling methods vary by stream type. One method is used in riffle/pool predominant streams, and the other method is for glide/pool predominant streams. For each stream type, macroinvertebrate sampling targets three habitats.

- For riffle/pool streams, the three habitats sampled are flowing water over coarse substrate, non-flowing water over depositional substrate, and rootmat substrate.
- For glide/pool streams, the three habitats sampled are non-flowing water over depositional substrate, large woody debris substrate, and rootmat substrate.

In some instances, one or more of the habitats sampled can be limited or missing from a stream reach, which may affect an MSCI score. Macroinvertebrate samples based on only two habitats may have an MSCI score equal to or greater than 16, but it is also possible that a missing habitat may lead to a decreased MSCI score. Although MDNR stream habitat assessment procedures take into account a number of physical habitat parameters from the sample reach (for example, riparian vegetation width, channel alteration, bank stability, bank vegetation protection, etc.), they do not exclusively measure the quality or quantity of the three predominant habitats from each stream. When evaluating potentially impaired macroinvertebrate communities, the number of habitats sampled, in addition to the stream habitat assessment score, will be considered to ensure MSCI scores less than 16 are properly attributed to poor water quality or poor/inadequate habitat condition.

Biologists responsible for conducting biological assessments will determine the extent to which habitat availability is responsible for a non-supporting (<16) MSCI score. If it is apparent that a non-supporting MSCI score was due to limited habitat, these effects will be stated in the biological assessment report. This limitation will then be considered when deciding which Listing Methodology category is most appropriate for an individual stream. This procedure, as part of an MDNR biological assessment, will aid in determining whether impaired macroinvertebrate samples have MSCI scores based on poor water quality conditions versus habitat limitations.

To ensure assessments are based on representative macroinvertebrate samples, samples collected during or shortly after prolonged drought, shortly after major flood events, or any other conditions that fall outside the range of environmental conditions under which reference streams in the EDU were sampled, will not be used to make an attainment decision for a Section 303(d) listing or any other water quality assessment purposes.

Sample “representativeness” is judged by Water Protection Program (WPP) staff after reading the biomonitoring report for that stream, and if needed, consultation with biologists from the department’s Environmental Services Program. Regarding smaller deviations from “normal” conditions, roughly 20 percent of reference samples failing to meet a fully biologically supporting MSCI score were collected following weather/climate extremes; as a result, biological criteria for a given EDU are inclusive of samples collected during not only ideal macroinvertebrate-rearing conditions, but also during the weather extremes that Missouri experiences.

#### ***Assessing Small Streams***

Occasionally, macroinvertebrate monitoring is needed to assess streams smaller than typical the wadeable/perennial reference streams listed in Table I of Missouri’s Water Quality Standards. Smaller streams may include Class C streams (streams that may cease flow in dry periods but maintain permanent pools which support aquatic life) or those that are unclassified. Assessing small streams involves comparing test stream and candidate reference stream MSCI scores first, to Wadeable/Perennial Reference Stream (WPRS) criteria, and second to each other. In MDNR’s Biological Criteria Database, there are 16 candidate reference streams labeled as Class P, 23 labeled as Class C, and 24 labeled as Class U. In previous work by MDNR, when the MSCI was calculated according to WPRS criteria, the failure rate for such candidate reference streams was 31% for Class P, 39% for Class C, and 70% for Class U. The data trend showed a higher failure rate for increasingly smaller high quality streams when scored using WPRS biological criteria. This trend demonstrates the need to include the utilization of candidate reference streams in biological stream assessments.

For test streams that are smaller than wadeable perennial reference streams, MDNR also samples five candidate reference streams (small control streams) of same or similar size and Valley Segment Type (VST) in the same EDU twice during the same year the test stream is sampled (additional information about the selection small control streams is provided below). Although in most cases the MDNR samples small candidate reference streams concurrently with test streams, existing data may be used if a robust candidate reference stream data set exists for the EDU.

If the ten small candidate reference stream scores are similar to wadeable perennial reference stream criteria, then they and the test stream are considered to have a Class C or Class P general warm water beneficial use, and the MSCI scoring system in the LMD should be used. If the small candidate reference streams have scores lower than the wadeable perennial reference streams, the assumption is that the small candidate reference streams, and the test stream, represent designated uses related to stream size that are not yet approved by EPA in the state’s water quality standards. The current assessment method for test streams that are smaller than reference streams is stated below.

- If the ten candidate reference stream (small control stream) scores are similar to WPRSs and meet LMD criteria for an unimpaired macroinvertebrate community, then the test stream will be assessed using MSCI based procedures in the LMD.
- If the ten candidate reference stream scores are lower than those of WPRSs and do not meet the LMD criteria for an unimpaired macroinvertebrate community, then:
  - a) The test stream will be assessed as having an unimpaired macroinvertebrate community if the test stream scores meet the LMD criteria for an unimpaired community;
  - b) The test stream data will be judged inconclusive if test stream scores are similar to candidate reference stream scores;
  - c) The test stream will be assessed as having a “suspect” macroinvertebrate community if its scores are found to be low but statistically close to candidate reference streams; or,
  - d) The test stream will be assessed as having an “impaired” macroinvertebrate community if its scores are found to be statistically lower than the candidate reference streams.

This method of assessing small streams will be used only until such time as the aquatic habitat protection use categories based on watershed size classifications of Headwater, Creek, Small River, Large River and Great River are promulgated into Missouri Water Quality Standards and appropriate biological metrics are established for stream size and permanence.

The approach for determining a “suspect” or “impaired” macroinvertebrate community will be made using a direct comparison between all streams being evaluated, which may include the use of percent and/or mean calculations as determined on a case by case basis. All work will be documented on the macroinvertebrate assessment worksheet and be made available during the public notice period.

#### ***Selecting Small Candidate Reference Streams***

Accurately assessing streams that are smaller than reference streams begins with properly selecting small candidate reference streams. Candidate reference streams are smaller than WPRS streams and have been identified as “best available” reference stream segments in the same EDU as the test stream according to watershed, riparian, and in-channel conditions. The selection of candidate reference streams is consistent with framework provided by Hughes *et al.* (1986) with added requirements that candidate reference streams must be from the same EDU and have the same or similar values for VST parameters. If candidate reference streams perform well when compared to WPRS, then test streams of similar size and VST are expected to do so as well. VST parameters important for selection are based on temperature, stream size, flow, geology, and relative gradient, with emphasis placed on the first three parameters.

The stepwise process for candidate reference stream selection is listed below.

1. Determine test stream reaches to be assessed.
2. Identify appropriate EDU.
3. Determine five variable VST of test stream segments (1<sup>st</sup> digit = temperature; 2<sup>nd</sup> digit = size; 3<sup>rd</sup> digit = flow; 4<sup>th</sup> digit = geology; and 5<sup>th</sup> digit = relative gradient).
4. Filter all stream segments within the same EDU for the relevant five variable VSTs (1<sup>st</sup> and 2<sup>nd</sup> digits especially critical for small streams).
5. Filter all potential VST stream segments for stressors against available GIS layers (e.g. point source, landfills, CAFOs, lakes, reservoirs, mining, etc.).
6. Filter all potential VST stream segments against historical reports and databases.
7. Develop candidate stream list with coordinates for field verification.
8. Field verify candidate list for actual use (e.g. animal grazing, in-stream habitat, riparian habitat, migration barriers, representativeness, gravel mining, and other obvious human stressors).
9. Rank order candidate sites, eliminate obvious stressed sites, and select at least top five sites.
10. Calculate land use-land cover and compare to EDU.
11. Collect chemical, biological, habitat, and possibly sediment field data.
12. After multiple sampling events evaluate field data, land use, and historical data in biological assessment report.
13. If field data are satisfactory, retain candidate reference stream label in database.

#### *Fish Community Data*

The department utilizes fish community data to determine if aquatic life use is supported in certain types of Missouri streams. When properly evaluated, fish communities serve as important indicators of stream health. In Missouri, fish communities are surveyed by the MDC. MDC selects an aquatic subregion to sample each year, and therein, surveys randomly selected streams of 2<sup>nd</sup> to 5<sup>th</sup> order in size. Fish sampling follows procedures described in the document *Resource Assessment and Monitoring Program: Standard Operational Procedures—Fish Sampling* (Combes 2011). Numeric biocriteria for fish are represented by the fish Index of Biotic Integrity (fIBI). Development of the fIBI is described in the document *Biological Criteria for Stream Fish Communities of Missouri* (Doisy et al. 2008).

The fIBI is a multi-metric index made up of nine individual metrics, which include:

- number (#) of -native individuals;
- # of native darter species;

- # of native benthic species;
- # of native water column species;
- # of native minnow species;
- # of all native lithophilic species;
- percentage (%) of native insectivore cyprinid individuals;
- % of native sunfish individuals; and,
- % of the three top dominant species.

Values for each metric, as directly calculated from the fish community sample, are converted to unitless scores of 1, 3, or 5 according to criteria in Doisy *et al.* (2008). The fIBI is then calculated by adding these unitless values together for a total possible score of 45. Doisy *et al.* (2008) established an impairment threshold of 36 (where the 25<sup>th</sup> percentile of reference sites represented a score of 37), with values equal to or greater than 36 representing unimpaired communities, and values less than 36 representing impaired communities. For more information regarding fIBI scoring, please see Doisy *et al.* (2008).

Based on consultation between the department and MDC, the fIBI impairment threshold value of 36 was used as the numeric biocriterion translator for making an attainment decision for aquatic life ([Appendix C](#)). Work by Doisy *et al.* (2008) focused on streams 3<sup>rd</sup> to 5<sup>th</sup> order in size, and the fIBI was only validated for streams in the Ozark ecoregion, not for streams in the Central Plains and Mississippi Alluvial Basin. Therefore, when assessing streams with the fIBI, the index may only be applied to streams 3<sup>rd</sup> to 5<sup>th</sup> order in size from the Ozark ecoregion. Assessment procedures are outlined below.

Comment [R22]: Previously Table 1.2

#### Full Attainment

- For seven or fewer samples and following MDC RAM fish community protocols, 75% of fIBI scores must be 36 or greater. Fauna achieving these scores are considered to be very similar to Ozark reference streams.
- For eight or more samples, the percent of samples scoring 36 or greater must be statistically similar to representative reference or control streams. To determine statistical similarity, a binomial probability Type I error rate (0.1) is calculated based on the null hypothesis that the test stream would have the same percentage (75%) of fIBI scores greater than 36 as reference streams. If the Type I error rate is more than the significance level  $\alpha=0.1$ , the fish community would be rated as unimpaired.

#### Non-Attainment

- For seven or fewer samples and following MDC RAM fish community protocols, 75% of the fIBI scores must be lower than 36. Fauna achieving these scores are considered to be substantially different than regional reference streams.

- For eight or more samples, the percent of samples scoring 36 or less must be statistically dissimilar to representative reference or control streams. To determine statistical dissimilarity, a binomial probability Type I error rate is calculated based on the null hypothesis that the test stream would have the same percentage (75%) of fIBI scores greater than 36 as reference streams. If the Type I error rate is less than 0.1, the null hypothesis is rejected and the fish community would be rated as impaired.

**Data will be judged inconclusive when outcomes do not meet requirements for decisions of full or non-attainment.**

With the exception of two subtle differences, use of the binomial probability for fish community samples will follow the example provided for macroinvertebrate samples in the previous section. First, instead of test stream samples being compared to reference streams of the same EDU, they will be compared to reference streams from the Ozark ecoregion. Secondly, the probability of success used in the binomial distribution equation will always be set to 0.70 since Appendix D states to “rate a stream as impaired if biological criteria reference stream frequency of fully biologically supporting scores is greater than five percent more than the test stream.”

Although 1<sup>st</sup> and 2<sup>nd</sup> order stream data will not be used to judge a stream as impaired for Section 303(d) purposes, the department may use the above assessment procedures to judge 1<sup>st</sup> and 2<sup>nd</sup> order streams as unimpaired. Moreover, should samples contain fIBI scores less than 29, the department may judge the stream as “suspected of impairment” using the above procedures.

***Considerations for the Influence of Habitat Quality and Sample Representativeness***

Low fIBI scores that are substantially different than reference streams could be the result of water quality problems, habitat problems, or both. When low fIBI scores are established, it is necessary to review additional information to differentiate between an impairment caused by water quality and one that is caused by habitat. The collection of a fish community sample is also accompanied by a survey of physical habitat from the sampled reach. MDC sampling protocol for stream habitat follows procedures provided by Peck *et al.* (2006). With MDC guidance, the department utilizes this habitat data and other available information to assure that an assessment of aquatic life attainment based on fish data is only the result of water quality, and that an impairment resulting from habitat is categorized as such. This section describes the procedures used to assure low fIBI scores are the result of water quality problems and not habitat degradation. The information below outlines the department’s provisional method to identify unrepresentative samples and low fIBI scores with questionable habitat condition, and ensure corresponding fish IBI scores are not used for Section 303(d) listing.

Comment [R23]: Previously B-1

- a) Following recommendations from the biocriteria workgroup, the department will consult MDC about the habitat condition of particular streams when assessing low fIBI scores.
- b) Samples may be considered for Section 303(d) listing ONLY if they were collected in the Ozark ecoregion, and the samples were collected during normal representative conditions, based upon best professional judgment from MDC staff. Samples collected from the Central Plains and Mississippi Alluvial Basin are excluded from Section 303(d) listing.
- c) Only samples from streams 3rd to 5th order in size may be considered for Section 303(d) listing. Samples from 1st or 2nd order stream sizes are excluded from Section 303(d) consideration; however, they may be placed into Categories 2B and 3B if impairment is suspected, or into Categories 1, 2A, or 3A if sample scores indicate a stream is unimpaired. Samples from lower stream orders are surveyed under a different RAM Program protocol than 3rd to 5th order streams.
- d) Samples that are ineligible for Section 303(d) listing include those collected from losing streams, as defined by the Department of Geology and Land Survey, or; collected in close proximity to losing streams. Additionally, ineligible samples may include those collected on streams that were considered to have natural flow issues (such as streams reduced predominately to subsurface flow) preventing good fish IBI scores from being obtained, as determined through best professional judgment of MDC staff.
- e) Fish IBI scores must be accompanied by habitat samples with a QCPH1 habitat index score. MDC was asked to analyze meaningful habitat metrics and identify samples where habitat metrics seemed to indicate potential habitat concerns. As a result, a provisional index named QCPH1 was developed. QCPH1 values less than 0.39 indicate poor habitat, and values greater than 0.39 suggest adequate habitat is available. The QCPH1 comprises six sub-metrics indicative of substrate quality, channel disturbance, channel volume, channel spatial complexity, fish cover, and tractive force and velocity.

The QCPH1 index is calculated as follows:

$$QCPH1 = ((\text{Substrate Quality} * \text{Channel Disturbance} * \text{Channel Volume} * \text{Channel Spatial Complexity} * \text{Fish Cover} * \text{Tractive Force \& Velocity})^{1/6})$$

Where sub-metrics are determined by:

$$\text{Substrate Quality} = [(\text{embeddedness} + \text{small particles})/2] * [(\text{filamentous algae} + \text{aquatic macrophyte})/2] * \text{bedrock and hardpan}$$

**Channel Disturbance** = concrete \* riprap \* inlet/outlet pipes \*  
relative bed stability \* residual pool observed to expected ratio

**Channel Volume** = [(dry substrate+width depth product + residual  
pool + wetted width)/4]

**Channel Spatial Complexity** = (coefficient of variation of mean  
depth + coefficient of variation of mean wetted width + fish cover  
variety)/3

**Fish Cover** = [(all natural fish cover + ((brush and overhanging  
vegetation + boulders + undercut bank + large woody debris)/4) +  
large types of fish cover)/3]

**Tractive Force & Velocity** = [(mean slope + depth \* slope)/2]

Unimpaired fish IBI samples (fIBI  $\geq 36$ ) with QCPHI index scores below the 0.39 threshold value, or samples without a QCPHI score altogether, are eliminated from consideration for Category 5 and instead placed into Categories 2B or 3B should an impairment be suspected. Impaired fish communities (fIBI  $< 36$ ) with QCPHI scores  $< 0.39$  can be placed into Category 4C (non-discrete pollutant/habitat impairment). Impaired fish communities (fIBI  $< 36$ ) with adequate habitat scores (QCPHI  $> 0.39$ ) can be placed into Category 5. Appropriate streams with unimpaired fish communities and adequate habitat (QCPHI  $> 0.39$ ) may be used to judge a stream as unimpaired.

Similar to macroinvertebrates, assessment of fish community information must be based on data coded level three or four as described in Section II.C of [this document](#). Data coded as levels three and four represent environmental data with the greatest degree of assurance, and thus, assessments will include multiple samples from a single site, or samples from multiple sites within a single reach.

Following the department's provisional methodology, fish community samples available for assessment (using procedures in [Appendix C & D](#)) include only those from 3rd to 5th order Ozark Plateau streams, collected under normal, representative conditions, where habitat seemed to be good, and where there were no issues with inadequate flow or water volume.

- Other Biological Data

On a case by case basis, the department may use biological data other than MSCI or fIBI scores for assessing attainment of aquatic life. Other biological data may include information on single indicator aquatic species that are ecologically or recreationally important, or individual measures of community health that respond predictably to

Comment [R24]: Previously Table 1.2, B-1 & B-2

environmental stress. Measures of community health could be represented by aspects of structure, composition, individual health, and processes of the aquatic biota. Examples could include measures of density or diversity of aquatic organisms, replacement of pollution intolerant taxa, or even the presence of biochemical markers.

Other biological data should be collected under a well vetted study that is documented in a scientific report, a weight of evidence- approach should be established, and the report should be referenced in the 303(d) listing worksheet. If other biological data is a critical component of the community and has been adversely affected by the presence of a pollutant or stressor, then such data would indicate a water body is impaired. The department's use of other biological data is consistent with EPA's policy on independent applicability for making attainment decisions, which is intended to protect against dismissing valuable information when diagnosing an impairment of aquatic life.

The use of other biological data in water body assessments occurs infrequently, but when available, it is usually assessed in combination with other information collected within the water body of interest. The department will avoid using other biological data as the sole justification for a Section 303(d) listing; however, other biological data will be used as part of a weight of evidence analysis for making the most informed assessment decision.

- Toxic Chemicals

- Water

- For the interpretation of toxicity test data, standard acute or chronic bioassay procedures using freshwater aquatic fauna such as, but not limited to, *Ceriodaphnia dubia*, Fathead Minnows (*Pimephales promelas*), *Hyalella azteca*, or Rainbow Trout (*Oncorhynchus mykiss*)<sup>18</sup> will provide adequate evidence of toxicity for 303(d) listing purposes. Microtox<sup>®</sup> toxicity tests may be used to list a water as affected by "toxicity" only if there are data of another kind (freshwater toxicity tests, sediment chemistry, water chemistry, or biological sampling) that indicate water quality impairment.

- For any given water, available data may occur throughout the system and/or be concentrated in certain areas. When the location of pollution sources are known, the department reserves the right to assess data representative of impacted conditions separately from data representative of unimpacted conditions. Pollution sources include those that may occur at discrete points along a water body, or those that are more diffuse.

- Sediment

- For toxic chemicals occurring in benthic sediments, data interpretation will include calculation of a geometric mean for specific toxins from an adequate number of samples, and comparing that value to a corresponding Probable Effect Concentration (PEC) given by MacDonald *et al.* (2000). The PEC is the level of a pollutant above which harmful effects

<sup>18</sup> Reference 10 CSR 20-7.015(9)(L) for additional information

on the aquatic community are likely to be observed. –MacDonald (2000) gave an estimate of accuracy for the ability of individual PECs to predict toxicity. For all metals except arsenic, pollutant geometric means will be compared to 150% of the recommended PEC values. This comparison should meet confidence requirements applied elsewhere in [this document](#). When multiple contaminants occur in sediment, toxicity may occur even though the level of each individual pollutant does not reach toxic levels. The method of estimating the synergistic effects of multiple pollutants in sediments [is described below](#).

#### ***The Meaning of the Sediment Quotient and How to Calculate It***

[Although](#) sediment criteria in the form of a PEC are given for several individual contaminants, it is recognized that when multiple contaminants occur in sediment, toxicity may occur even though the level of each individual pollutant does not reach toxic levels. The method of estimating the synergistic effects of multiple pollutants in sediments given in MacDonald *et al.* (2000) includes the calculation of a [PECQ](#). PECQs greater than 0.75 will be judged as toxic.

This calculation is made by dividing the pollutant concentration in the sample by the PEC value for that pollutant. For single samples, the quotients are summed, and then normalized by dividing that sum by the number of pollutants in the formula. When multiple samples are available, the geomean (as calculated for specific pollutants) will be placed in the numerator position for each pollutant included in the equation.

Example:- A sediment sample contains the following results in mg/kg:

Arsenic 2.5, Cadmium 4.5, Copper 17, Lead 100, [and](#) Zinc 260.

The PEC values for these five pollutants in respective order are:

33, 4.98, 149, 128, [and](#) 459 [mg/kg](#).

PECQ =

$$[(2.5/33) + (4.5/4.98) + (17/149) + (100/128) + (260/459)]/5 = 0.488$$

#### ***Using PECQ to Judge Toxicity***

Based on research by MacDonald *et al.* (2000) 83% of sediment samples with PECQ less than 0.5 were non-toxic while 85% of sediment samples with PECQ greater than 0.5 were toxic. Therefore, to accurately assess the synergistic effects of sediment contaminants on aquatic life, the department will judge PECQ greater than 0.75 as toxic.

- **Duration of Assessment Period**

Except where the assessment period is specifically noted in [Appendix B](#), the time period [during](#) which data will be used in making the assessments will be determined by data age and

**Comment [D25]:** Moved the previous Appendix D here

**Comment [R26]:** Previously Table I.1

data code considerations, as well as representativeness considerations such as those described in footnote 14.

- Assessment of Tier Three Waters

Waters given Tier Three protection by the antidegradation rule at 10 CSR 20-7.031(2), shall be considered impaired if data indicate water quality has been reduced in comparison to its historical quality. Historical quality is determined from past data that best describes a water body's water quality following promulgation of the antidegradation rule and at the time the water was given Tier Three protection.

Historical data gathered at the time waters were given Tier Three protection will be used if available. Because historical data may be limited, the historical quality of the waters may be determined by comparing data from the assessed segment with data from a "representative" segment. A representative segment is a body or stretch of water that best reflects the conditions that probably existed at the time the antidegradation rule first applied to the waters being assessed. Examples of possible representative data include 1) data from stream segments upstream of assessed segments that receive discharges, and 2) data from other water bodies in the same ecoregion having similar watershed and landscape characters. These representative stream segments also would be characterized by receiving discharges similar to the quality and quantity of historic discharges of the assessed segment. The assessment may also use data from the assessed segment gathered between the time of the initiation of Tier Three protection and the last known time in which upstream discharges, runoff, and watershed conditions remained the same, provided that the data do not show any significant trends of declining water quality during that period.

**Comment [D27]:** rearranged wording to provide clarity.

The data used in the comparisons will be tested for normality and an appropriate statistical test will be applied. The null hypothesis for statistical analysis will be that water quality at the test segment and representative segment is the same. This will be a one-tailed test (the test will consider only the possibility that the assessed segment has poorer water quality) with the alpha level of 0.1, meaning that the test must show greater than a 90 percent probability that the assessed segment has poorer water quality than the representative segment before the assessed segment can be listed as impaired.

- Other Types of Information

1. Observation and evaluation of waters for noncompliance with state narrative water quality criteria. Missouri's narrative water quality criteria, as described in 10 CSR 20-7.031 Section (3), may be used to evaluate waters when a quantitative (narrative) value can be applied to the pollutant. These narrative criteria apply to both classified and unclassified waters and prohibit the following in waters of the state:

- a. Waters shall be free from substances in sufficient amounts to cause the formation of putrescent, unsightly, or harmful bottom deposits or prevent full maintenance of beneficial uses;

**Comment [D28]:** Replaced the summarized wording with the wording stated in WQS.

- b. Waters shall be free from oil, scum, and floating debris in sufficient amounts to be unsightly or prevent full maintenance of beneficial uses;
- c. Waters shall be free from substances in sufficient amounts to cause unsightly color or turbidity, offensive odor, or prevent full maintenance of beneficial uses;
- d. Waters shall be free from substances or conditions in sufficient amounts to result in toxicity to human, animal, or aquatic life;
- e. There shall be no significant human health hazard from incidental contact with the water;
- f. There shall be no acute toxicity to livestock or wildlife watering;
- g. Waters shall be free from physical, chemical, or hydrologic changes that would impair the natural biological community;
- h. Waters shall be free from used tires, car bodies, appliances, demolition debris, used vehicles or equipment, and solid waste as defined in Missouri's Solid Waste Law, section 260.200, RSMo, except as the use of such materials is specifically permitted pursuant to sections 260.200–260.247, RSMo;

2. Habitat assessment protocols for wadeable streams have been established and are conducted in conjunction with sampling aquatic macroinvertebrates and fish. Methods for evaluating aquatic macroinvertebrate and fish community data include assessment procedures that account for the presence or absence of representative habitat quality. The department will not use habitat data alone for assessment purposes.

#### **E. Other 303(d) Listing Considerations**

- Adding to the Existing List or Expanding the Scope of Impairment to a Previously Listed Water,

The listed portion of impaired water bodies may be increased based on recent monitoring data following the guidelines in this document. One or more new pollutants may be added to the listing for a water body already on the list based on recent monitoring data following these same guidelines. Waters not previously listed may be added to the list following the guidelines in this document.

- Deleting from the Existing List or Decreasing the Scope of Impairment to a Previously Listed Water

The listed portion of an impaired water body may be decreased based on recent monitoring data following the guidelines in this document. One or more pollutants may be deleted from the listing for a water body already on the list based on recent monitoring data following guidelines in Appendix D. Waters may be completely removed from the

list for several reasons<sup>19</sup>, the most common being (1) water has returned to compliance with water quality standards, or (2) the water has an approved TMDL study or Permit in Lieu of a TMDL.

#### F. Prioritization of Waters for TMDL Development

Section 303(d) of the Clean Water Act and federal regulation 40 CFR 130.7(b)(4) require states to submit a priority ranking of waters requiring TMDLs. The department will prioritize development of TMDLs based on several variables including:

- social impact/public interest and risk to public health
- complexity and cost (including consideration of budget constraints), availability of data of sufficient quality and quantity for -TMDL modeling
- court orders, consent decrees, or other formal agreements
- source of impairments
- existence of appropriate numeric quality criteria, and
- implementation potential and amenability of the problem to treatment.

The department's TMDL schedule will represent its prioritization. The TMDL Program develops the TMDL schedule and maintains it at the following website:  
<http://www.dnr.mo.gov/env/wpp/tmdl/>.

#### 4. Resolution of Interstate/International Disagreements

The department will review the draft 303(d) Lists of all other states with which it shares a border (Missouri River, Mississippi River, Des Moines River and the St. Francis River) or other interstate waters. Where the listing for the same water body in another state is different than the one in Missouri, the department will request the data and the listing justification. These data will be reviewed following the evaluation guidelines in this document. The Missouri Section 303(d) list may be changed pending the evaluation of this additional data.

#### G. Statistical Considerations

The most recent EPA guidance on the use of statistics in the 303(d) listing methodology document is given in Appendix A. Within this guidance there are three major recommendations regarding statistics:

- Provide a description of analytical tools the state uses under various circumstances;
- When conducting hypothesis testing, explain the various circumstances under which the burden of proof is placed on proving the water is impaired and when it is placed on proving the water is unimpaired, and
- Explain the level of statistical significance ( $\alpha$ ) used under various circumstances.

<sup>19</sup> See, "Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act". USEPA, Office of Water, Washington DC.

- Description of Analytical Tools

Appendix D, describes the analytical tools the department will use to determine whether a water body is impaired and whether or when a listed water body is no longer impaired.

- Rationale for the Burden-of-Proof

Hypothesis testing is a common statistical practice. The procedure involves first stating a hypothesis you want to test, such as “the most frequently seen color on clothing at a St. Louis Cardinals game is red” and then the opposite or null hypothesis “red is not the most frequently seen color on clothing at a Cardinals game.” Then a statistical test is applied to the data (a sample of the predominant color of clothing worn by 200 fans at a Cardinals game on July 12) and based on an analysis of that data, one of the two hypotheses is chosen as correct.

In hypothesis testing, the burden-of-proof is always on the alternate hypothesis. In other words, there must be very convincing data to make us conclude that the null hypothesis is not true and that we must accept the alternate hypothesis. How convincing the data must be is stated as the “significance level” of the test. A significance level of  $\alpha=0.10$  means that there must be at least a 90 percent probability that the alternate hypothesis is true before we can accept it and reject the null hypothesis.

For analysis of a specific kind of data, either the test significance level or the statement of null and alternative hypotheses, or both, can be varied to achieve the desired degree of statistical rigor. The department has chosen to maintain a consistent set of null and alternate hypotheses for all our statistical procedures. The null hypothesis will be that the water body in question is unimpaired and the alternate hypothesis will be that it is impaired. Varying the level of statistical rigor will be accomplished by varying the test significance level. For determining impairment (Appendix D) test significance levels are set at either  $\alpha=0.1$  or  $\alpha=0.4$ , meaning the data must show at minimum 90% or 60% probability, respectively, that the water body is impaired. However, if the department retained these same test significance levels in determining when an impaired water body had been restored to an unimpaired status (Appendix D) some undesirable results can occur.

**Comment [D29]:** Previously Table B-2

For example, using a 0.1 significance level for determining both impairment and non-impairment, if the sample data indicate the stream had a 92 percent probability of being impaired, it would be rated as impaired. If subsequent data were collected and added to the database, and the data now showed the water had an 88 percent chance of being impaired, it would be rated as unimpaired. Judging as unimpaired a water body with only a 12 percent probability of being unimpaired is clearly a poor decision. To correct this problem, the department will use a test significance level of 0.4 for some analytes and 0.6 for others. This will increase our confidence in determining compliance with criteria to 40 percent and 60 percent, respectively under the worst case conditions, and for most databases will provide an even higher level of confidence.

- Level of Significance Used in Tests

The choice of significance levels is largely related to two concerns. The first concern is with matching error rates with the severity of the consequences of making a decision error. The second addresses the need to balance, to the degree practicable, Type I and Type II error rates. For relatively small number of samples, the disparity between Type I and Type II errors can be large. The tables 2.0 and 3.0 below shows error rates calculated using the binomial distribution for two very similar situations. Type I error rates are based on a stream with a 10 percent exceedence rate of a standard, and Type II error rates are based on a stream with a 15 percent exceedence rate of a standard. Note that when sample size remains the same, Type II error rates increase as Type I error rates decrease (Table 2.0). Also note that for a given Type I error rate, the Type II error rate declines as sample size increases (Table 3.0).

Comment [R30]: Previously B-3

**Table 2.0.** Effects of Type I error rates on Type II error rates. Type I error rates are based on a stream with a 10 percent exceedence rate of a standard and Type II error rates for a stream with a 15 percent exceedence rate of a standard.

Comment [R31]: Previously Table B-3

Total No. of Samples	No. Samples Meeting Std.	Type I Error Rate	Type II Error Rate
18	17	0.850	0.479
18	16	0.550	0.719
18	15	0.266	0.897
18	14	0.098	0.958
18	13	0.028	0.988

**Table 3.0.** Effects of Type I error rates and sample size on Type II error rates. Type I error rates are based on a stream with a 10 percent exceedence rate of a standard and Type II error rates for a stream with a 15 percent exceedence rate of a standard.

Comment [R32]: Previously Table B-4

Total No. of Samples	No. Samples Meeting Std.	Type I Error Rate	Type II Error Rate
6	5	0.469	0.953
11	9	0.303	0.930
18	15	0.266	0.897
25	21	0.236	0.836

- Use of the Binomial Probability Distribution for Interpretation of the 10 Percent Rule

There are two options for assessing data for compliance with the 10 percent rule. One is to simply calculate the percent of time the criterion value is not met, and to judge the water to be impaired if this value is greater than 10 percent. The second method is to use some evaluative procedure that can review the data and provide a probability statement regarding compliance with the 10 percent rule. Since the latter option allows assessment decisions relative to specific test significance levels

and the first option does not, the latter option is preferred. The procedure chosen is the binomial probability distribution and calculation of the Type I error rate.

- Other Statistical Considerations

Prior to calculation of confidence limits, the normality of the data set will be evaluated. If normality is improved by a data transformation, the confidence limits will be calculated on the transformed data.

Time of sample collection may be biased and interfere with an accurate measurement of frequency of exceedance of a criterion. Data sets composed mainly or entirely of storm water data or data collected only during a season when water quality problems are expected could result in a biased estimate of the true exceedance frequency. In these cases, the department may use methods to estimate the true annual frequency and display these calculations whenever they result in a change in the impairment status of a water body.

For waters judged to be impaired based on biological data where data evaluation procedures are not specifically noted in Table 1, the statistical procedure used, test assumptions, and results will be reported.

- Examples of Statistical Procedures

*Two Sample "t" Test for Color*

Null Hypothesis: Amount of color is no greater in a test stream than in a control stream. As stated, this is a one-sided test, meaning that we are only interested in determining whether or not the color level in the test stream is greater than in a control stream. If the null hypothesis had been "amount of color is different in the test and control streams," we would have been interested in determining if the amount of color was either less than or greater than the control stream, a two-sided test.

Significance Level:  $\alpha=0.10$

Data Set: Platinum-Cobalt color units data for the test stream and a control stream samples collected at each stream on same date.

Test Stream	70	45	35	45	60	60	80
Control Stream	50	40	20	40	30	40	75
Difference (T-C)	20	5	15	5	30	20	5

Statistics for the Difference: Mean = 14.28, standard deviation = 9.76,  $n = 7$

Calculated "t" value = (square root of n)(mean)/standard deviation = 3.86

Tabular "t" value is taken from a table of the "t" distribution for 2 alpha (0.20) and n-1 degrees of freedom. Tabular "t" = 1.44.

**Comment [D33]:** Removed from sentence "for data sets up to size 30. Use of the binomial probability is difficult for larger sample sizes. And for these larger data sets impairment will be determined by making direct comparison of percent of samples not compliant with the criterion value with the ten percent guideline."

Since calculated "t" value is greater than tabular t value, reject the null hypothesis and conclude that the test stream is impaired by color.

Statistical Procedure for Mercury in Fish Tissue

Data Set: data in  $\mu\text{g/Kg}$  130, 230, 450. Mean = 270, Standard Deviation = 163.7

The 60% Lower Confidence Limit Interval = the sample mean minus the quantity:

$$((0.253)(163.7)/\text{square root } 3) = 23.9. \text{ -Thus the 60\% LCL Confidence Interval is } 246.088 \mu\text{g/Kg.}$$

The criterion value is 300  $\mu\text{g/Kg}$ . Therefore, since the 60% LCL Confidence Interval is less than the criterion value, the water is judged to be unimpaired by mercury in fish tissue, and the water body is placed in either Category 2B or 3B.


## H. References

- Barbour, M.T., J. Gerritsen, G.E. Griffith, R. Frydenborg, E. McCarron, J.S. White, M.L. Bastian. 1996. A framework for biological criteria for Florida streams using benthic macroinvertebrates. *Journal of the North American Benthological Society* 15(2): 185-211.
- Doisy, K.E., C.F. Rabeni, M.D. Combes, and R.J. Sarver. 2008. Biological Criteria for Stream Fish Communities of Missouri. Final Report to the United States Environmental Protection Agency. Missouri Cooperative Fish and Wildlife Research Unit, Columbia, Missouri. 91 pp.
- Hughes, R.M., D.P. Larsen, and J.M. Omernik. 1986. Regional reference sites: a method for assessing stream pollution. *Environmental Management* 10(5): 625-629.
- Ohio Environmental Protection Agency. 1990. The Use of Biocriteria in the Ohio EPA Surface Water Monitoring and Assessment Program. Columbus, Ohio.
- Fischer, S. and M. Combes. 2011. Resource Assessment and Monitoring Program: Standard Operating Procedures – Fish Sampling. Missouri Department of Conservation, Jefferson City, Missouri.
- MacDonald, D.D, Ingersoll, C. G., Berger, T. A. *et al.* 2000. Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems. *Arch. Environ. Contamination Toxicology*. 39, 20-31.
- Missouri Department of Natural Resources. 2002. Biological Criteria for Wadeable/Perennial Streams of Missouri. Missouri Department of Natural Resources, Environmental Services Program, P.O. Box 176, Jefferson City, Missouri 65102. 32 pp.
- [Missouri Department of Natural Resources. 2010. Stream Habitat Assessment. Missouri Department of Natural Resources, Environmental Services Program, P.O. Box 176, Jefferson City, Missouri 65102. 40 pp.](#)
- Missouri Department of Natural Resources. 2012a. Semi-Quantitative Macroinvertebrate Stream Bioassessment. Missouri Department of Natural Resources, Environmental Services Program, P.O. Box 176, Jefferson City, Missouri 65102. 29 pp.
- Missouri Department of Natural Resources. 2014 (recertification). Taxonomic Levels for Macroinvertebrate Identifications. Division of Environmental Quality, Environmental Services Program, P.O. Box 176, Jefferson City, Missouri 65102. 39 pp.
- Peck, D.V., A.T. Herlihy, B.H. Hill, R.M. Hughes, P.R. Kaufmann, D.J. Klemm, J.M. Lazorchak, F.H. McCormick, S.A. Peterson, P.L. Ringold, T. Magee, and M.Cappaert. 2006. Environmental Monitoring and Assessment Program-Surface Waters Western Pilot Study: Field Operation Manual for Wadeable Streams. EPA/620/R-06/003. U.S.

Environmental Protection Agency, Office of Research and Development, Washington, D.C.

U.S. Environmental Protection Agency. 1996. Biological Criteria: Technical Guidance for Streams and Small Rivers. EPA 822-B-96-001. Office of Water, Washington D.C. 162 pp.

Proposed



## Appendix A

**Excerpt** from *Guidance for 2006 Assessment, Listing and Reporting Requirements Pursuant to Sections 303(d), 305(b) and 314 of the Clean Water Act*. July 29, 2005. USEPA pp. 39-41.

The document can be read in its entirety from the US. EPA web site:  
<http://water.epa.gov/lawsregs/lawsguidance/cwa/tmdl/upload/2006irg-report.pdf>

### G. How should statistical approaches be used in attainment determinations?

*The state's methodology should provide a rationale for any statistical interpretation of data for the purpose of making an assessment determination.*

#### *Description of statistical methods to be employed in various circumstances*

*The methodology should provide a clear explanation of which analytic tools the state uses and under which circumstances. EPA recommends that the methodology explain issues such as the selection of key sample statistics (arithmetic mean concentration, median concentration, or a percentile), null and alternative hypotheses, confidence intervals, and Type I and Type II error thresholds. The choice of a statistic tool should be based on the known or expected distribution of the concentration of the pollutant in the segment (e.g., normal or log normal) in both time and space.*

*Past EPA guidance (1997 305(b) and 2000 CALM) recommended making non-attainment decisions, for "conventional pollutants"<sup>20</sup> — TSS, pH, BOD, fecal coliform bacteria, and oil and grease<sup>13</sup> — when more than "10% of measurements exceed the water quality criterion." (However, EPA guidance has not encouraged use of the "10% rule" with other pollutants, including toxics.) Use of this rule when addressing conventional pollutants, is appropriate if its application is consistent with the manner in which applicable WQC are expressed. An example of a WQC for which an assessment based on the ten percent rule would be appropriate is the EPA acute WQC for fecal coliform bacteria, applicable to protection of water contact recreational use. This 1976-issued WQC was expressed as, "...no more than ten percent of the samples exceeding 400 CFU per 100 ml, during a 30-day period." Here, the assessment methodology is clearly reflective of the WQC.*

*On the other hand, use of the ten percent rule for interpreting water quality data is usually not consistent with WQC expressed either as: 1) instantaneous maxima not to be surpassed at any time, or 2) average concentrations over specified times. In the case of "instantaneous maxima (or minima) never to occur" criteria use of the ten percent rule typically leads to the belief that segment conditions are equal or better*

<sup>20</sup> There are a variety of definitions for the term "conventional pollutants." Wherever this term is referred to in this guidance, it means "a pollutant other than a toxic pollutant."

**Comment [D34]:** This entire section was updated with the exact wording from EPA's guidance document. A web link was also added for quick reference.

than specified by the WQC, when they in fact are considerably worse. (That is, pollutant concentrations are above the criterion-concentration a far greater proportion of the time than specified by the WQC.) Conversely, use of this decision rule in concert with WQC expressed as average concentrations over specific times can lead to concluding that segment conditions are worse than WQC, when in fact they are not.

If the state applies different decision rules for different types of pollutants (e.g., toxic, conventional, and non-conventional pollutants) and types of standards (e.g., acute vs. chronic criteria for aquatic life or human health), the state should provide a reasonable rationale supporting the choice of a particular statistical approach to each of its different sets of pollutants and types of standards.

1. Elucidation of policy choices embedded in selection of particular statistical approaches and use of certain assumptions EPA strongly encourages states to highlight policy decisions implicit in the statistical analysis that they have chosen to employ in various circumstances. For example, if hypothesis testing is used, the state should make its decision-making rules transparent by explaining why it chose either "meeting WQS" or "not meeting WQS" as the null hypothesis (rebuttable presumption) as a general rule for all waters, a category of waters, or an individual segment. Starting with the assumption that a water is "healthy" when employing hypothesis testing means that a segment will be identified as impaired, and placed in Category 4 or 5, only if substantial amounts of credible evidence exist to refute that presumption. By contrast, making the null hypothesis "WQS not being met" shifts the burden of proof to those who believe the segment is, in fact, meeting WQS.

Which "null hypothesis" a state selects could likely create contrasting incentives regarding support for additional ambient monitoring among different stakeholders. If the null hypothesis is "meeting standards," there were no previous data on the segment, and no additional existing and readily available data and information are collected, then the "null hypothesis" cannot be rejected, and the segment would not be placed in Category 4 or 5. In this situation, those concerned about possible adverse consequences of having a segment declared "impaired" might have little interest in collection of additional ambient data. Meanwhile, users of the segment would likely want to have the segment monitored, so they can be ensured that it is indeed capable of supporting the uses of concern. On the other hand, if the null hypothesis is changed to "segment not meeting WQS," then those that would prefer that a particular segment not be labeled "impaired" would probably want more data collected, in hopes of proving that the null hypothesis is not true.

Another key policy issue in hypothesis testing is what significance level to use in deciding whether to reject the null hypothesis. Picking a high level of significance for rejecting the null hypothesis means that great emphasis is being placed on avoiding a Type I error (rejecting the null hypothesis, when in fact, the null hypothesis is true). This means that if

*a 0.10 significance level is chosen, the state wants to keep the chance of making a Type I error at or below ten percent. Hence, if the chosen null hypothesis 2006 IR Guidance July 2005 41 is "segment meeting WQS," the state is trying to keep the chance of saying a segment is impaired – when in reality it is not – under ten percent.*

*An additional policy issue is the Type II errors (not rejecting the null hypothesis, when it should have been). The probability of Type II errors depends on several factors. One key factor is the number of samples available. With a fixed number of samples, as the probability of Type I error decreases, the probability of a Type II error increases. States would ideally collect enough samples so the chances of making Type I and Type II errors are simultaneously small. Unfortunately, resources needed to collect such numbers of samples are quite often not available.*

*The final example of a policy issue that a state should describe is the rationale for concentrating limited resources to support data collection and statistical analysis in segments where there are documented water quality problems or where the combination of nonpoint source loadings and point source discharges would indicate a strong potential for a water quality problem to exist.*

*EPA recommends that, when picking the decision rules and statistical methods to be utilized when interpreting data and information, states attempt to minimize the chances of making either of the two following errors:*

- Concluding the segment is impaired, when in fact it is not, and*
- Deciding not to declare a segment impaired, when it is in fact impaired.*

*States should specify in their methodology what significance level they have chosen to use, in various circumstances. The methodology would best describe in "plain English" the likelihood of deciding to list a segment that in reality is not impaired (Type I error if the null hypothesis is "segment not impaired"). Also, EPA encourages states to estimate, in their assessment databases, the probability of making a Type II error (not putting on the 303(d) list a segment that in fact fails to meet WQS), when: 1) commonly-available numbers of grab samples are available, and 2) the degree of variance in pollutant concentrations are at commonly encountered levels. For example, if an assessment is being performed with a WQC expressed as a 30-day average concentration of a certain pollutant, it would be useful to estimate the probability of a Type II error when the number of available samples over a 30 day period is equal to the average number of samples for that pollutant in segments state-wide, or in a given group of segments, assuming a degree of variance in levels of the pollutant often observed over typical 30 day periods.*

**Appendix B**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NUMERIC CRITERIA THAT ARE INCLUDED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

DESIGNATED USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>1</sup>	Notes
Overall use protection (all designated uses)	No data. Evaluated based on similar land use/ geology as stream with water quality data.	Not applicable	Given same rating as monitored stream with same land use and geology.	<b>Data Type Note:</b> This data type is used only for wide-scale assessments of aquatic biota and aquatic habitat for 305(b) Report purposes. This data type is not used in the development of the 303(d) List.
Any designated uses	No data available or where only effluent data is available. Results of dilution calculations or water quality modeling	Not applicable	Where models or other dilution calculations indicate noncompliance with allowable pollutant levels and frequencies noted in this table, waters may be added to Category 3B and considered high priority for water quality monitoring.	
Protection of Aquatic Life	Dissolved oxygen, water temperature, pH, total dissolved gases, oil and grease.	1-4	<b>Full:</b> No more than 10% of all samples exceed criterion. <b>Non-Attainment:</b> Requirements for full attainment not met.	<b>Compliance with Water Quality Standards Note:</b> Some sampling periods are wholly or predominantly during the critical period of the year when criteria violations occur. Where the monitoring program presents good evidence of a demarcation between seasons where criteria exceedences occur and seasons when they do not, the 10% exceedence rate will be based on an annual estimate of the frequency of exceedence.  <u>Chronic pH will be used in the LMD only if these criteria appear in the Code of State Regulations, and approved by the U.S. Environmental Protection Agency.</u>

**Comment [D35]:** Previously Table 1.1. Footnotes moved to "Notes" column

**Comment [D36]:** DO was combined with this row because it communicated the same information and footnote.

**Comment [D37]:** Previously 2016

**Appendix B**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NUMERIC CRITERIA THAT ARE INCLUDED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

DESIGNATED USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>1</sup>	Notes
Losing Streams	<i>E. coli</i> bacteria	1-4	<p><b>Full:</b> No more than 10% of all samples exceed criterion.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p> <p>The criterion for <i>E. coli</i> is 126 counts/100ml. 10 CSR 20-7.031 (4)(C)</p>	
Protection of Aquatic Life	Toxic chemicals	1-4	<p><b>Full:</b> No more than one acute toxic event in three years that results in a documented die-off of aquatic life such as fish, mussels, and crayfish (does not include die-offs due to natural origin). No more than one exceedence of acute or chronic criterion in the last three years for which data is available.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	
Protection of Aquatic Life	Nutrients in Lakes (total phosphorus, total nitrogen, plus chlorophyll)	1-4	<p><b>Full:</b> Nutrient levels do not exceed water quality standards following procedures stated in <a href="#">Appendix B</a>.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	<p><b>Compliance with Water Quality Standards Note:</b> Nutrient criteria will be used in the <del>2018</del> LMD only if these criteria appear in the Code of State Regulations, and approved by the U.S. Environmental Protection Agency.</p>
Human Health - Fish Consumption	Chemicals (water)	1-4	<p><b>Full:</b> Water quality does not exceed water quality standards following procedures stated in <a href="#">Appendix D</a>.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	

**Comment [R35]:** Previously Table 1.1. Footnotes moved to "Notes" column.

**Comment [D39]:** Previously 2016

**Comment [R38]:** Previously Table B-1

**Comment [D40]:** Previously Table B-1

**Appendix B**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NUMERIC CRITERIA THAT ARE INCLUDED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

DESIGNATED USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>1</sup>	Notes
Drinking Water Supply -Raw Water.	Chemical (toxics)	1-4	<p><b>Full:</b> Water Quality Standards not exceeded following procedures stated in <a href="#">Appendix I</a>.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	<p><b>Designated Use Note:</b> Raw water is water from a stream, lake or groundwater prior to treatment in a drinking water treatment plant.</p>
Drinking Water Supply- Raw Water	Chemical (sulfate, chloride, fluoride)	1-4	<p><b>Full:</b> <u>Water quality standards</u> not exceeded following procedures stated in <a href="#">Appendix I</a>.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	
Drinking Water Supply-Finished Water	Chemical (toxics)	1-4	<p><b>Full:</b> No Maximum Contaminant Level (MCL) violations based on Safe Drinking Water Act data evaluation procedures.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	<p><b>Compliance with Water Quality Standards Note:</b> Finished water data will not be used for analytes where water quality problems may be caused by the drinking water treatment process such as the formation of Trihalomethanes (THMs) or problems that may be caused by the distribution system (bacteria, lead, copper).</p>
Whole-Body-Contact Recreation and Secondary Contact Recreation	Fecal coliform or <i>E. coli</i> count	2-4	<p>Where there are at least five samples per year taken during the recreational season:</p> <p><b>Full:</b> <u>Water quality standards</u> not exceeded as a geometric mean, in any of the last three years for which data is available, for samples collected during seasons for which bacteria criteria apply.</p> <p><b>Non-Attainment:</b> Requirements for full attainment not met.</p>	<p><b>Compliance with Water Quality Standards Note:</b> A geometric mean of 206 cfu/100 ml for <i>E. coli</i> will be used as a criterion value for Category B Recreational Waters. Because Missouri's Fecal Coliform Standard ended December 31, 2008, any waters appearing on the 2008 303(d) List as a result of the Fecal Coliform Standard will be retained on the list with the pollutant listed as "bacteria" until sufficient <i>E. coli</i> sampling has determined the status of the water.</p>

**Comment [R35]:** Previously Table I.1. Footnotes moved to "Notes" column

**Comment [R41]:** Previously Table B-1

**Comment [R42]:** Previously Table B-1

**Comment [D43]:** This note was moved from the "Compliance with Water Quality Standards" column

**Appendix B**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NUMERIC CRITERIA THAT ARE INCLUDED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

DESIGNATED USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>1</sup>	Notes
Irrigation, Livestock and Wildlife Water	Chemical	1-4	Full: <u>Water quality standards</u> not exceeded following procedures stated in <a href="#">Appendix D</a> . Non-Attainment: Requirements for full attainment not met.	

<sup>1</sup> See section on Statistical Considerations, [Appendix C & D](#).

**Comment [R35]:** Previously Table 1.1. Footnotes moved to "Notes" column

**Comment [R44]:** Previously Table B-1

**Comment [D45]:** Previously Table B-1 & B-2

PROPOSAL

**Appendix C**

**Comment [R46]:** Previously Table 1.2  
 Footnotes moved into "Notes" column

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NARRATIVE CRITERIA BASED ON NUMERIC THRESHOLDS NOT CONTAINED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

BENEFICIAL USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>1</sup>	Notes
Overall use protection (all beneficial uses)	Narrative criteria for which quantifiable measurements can be made.	1-4	<p><b>Full:</b> Stream <b>condition</b> typical of reference or appropriate control streams in this region of the state.</p> <p><b>Non-Attainment:</b> The weight of evidence, based on the narrative criteria in 10 CSR 20-7.031(3), demonstrates the observed condition exceeds a numeric threshold necessary for the attainment of a beneficial use.</p> <p>For example:                      Color: Color as measured by the Platinum-Cobalt visual method (SM 2120 B) in a <b>water body</b> is statistically significantly higher than a control water.</p> <p>Objectionable Bottom Deposits: The bottom that is covered by sewage sludge, trash, or other materials reaching the water due to anthropogenic sources exceeds the amount in reference or control streams by more than <u>20</u> percent.</p> <p>Note: Waters in mixing zones and unclassified waters <b>that</b> support aquatic life on an intermittent basis shall be subject to acute toxicity criteria for protection of aquatic life. Waters in the initial Zone of Dilution shall not be subject to acute toxicity criteria.</p>	

**Comment [D47]:** Previously "apparent"

**Comment [R46]:** Previously Table 1.2  
 Footnotes moved into "Notes" column

**Appendix C**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NARRATIVE CRITERIA BASED ON NUMERIC THRESHOLDS NOT CONTAINED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

BENEFICIAL USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>ii</sup>	Notes
Protection of Aquatic Life	Toxic Chemicals	1-4	<p><u>Full</u>: No more than one acute toxic event in three years (does not include die-offs of aquatic life due to natural origin). No more than one exceedance of acute or chronic criterion in three years for all toxics.</p> <p><u>For hardness based metals with eight or fewer samples, the hardness value associated with the sample will be used to calculate the acute or chronic thresholds.</u></p> <p><u>For hardness based metals with more than eight samples, the reference percentile hardness provided in state water quality standards will be used to calculate the acute and chronic thresholds.</u></p> <p><u>Non-Attainment</u>: Requirements for full attainment not met.</p>	<p><b>Compliance with Water Quality Standards Note:</b> The test result must be representative of water quality for the entire time period for which acute or chronic criteria apply. For ammonia the chronic exposure period is 30 days, for all other toxics 96 hours. The acute exposure period for all toxics is 24 hours, except for ammonia which has a one hour exposure period. The department will review all appropriate data, including hydrographic data, to ensure only representative data is used. Except on large rivers where storm water flows may persist at relatively unvarying levels for several days, grab samples collected during storm water flows will not be used for assessing chronic toxicity criteria.</p> <p><b>Compliance with Water Quality Standards Note:</b> In the case of toxic chemicals occurring in toxic sediment rather than in water, the numeric thresholds used to determine the need for further evaluation will be the Probable Effect Concentrations proposed in "Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems" by MacDonald, D.D. <i>et al.</i> Arch. Environ. Contam. Toxicol. 39:20-31 (2000). These Probable Effect Concentrations are as follows: 33 mg/kg As; 4.98 mg/kg Cd; 111 mg/kg Cr; 149 mg/kg Cu; 48.6 mg/kg Ni; 128 mg/kg Pb; 459 mg/kg Zn; 561 µg/kg naphthalene; 1170 µg/kg phenanthrene; 1520 µg/kg pyrene; 1050 µg/kg benzo(a)anthracene; 1290 µg/kg chrysene; 1450 µg/kg benzo(a)pyrene; 22,800 µg/kg total polycyclic aromatic hydrocarbons; 676 µg/kg total PCBs; chlordane 17.6 ug/kg; Sum DDE 31.3 ug/kg; lindane (gamma-BHC) 4.90 ug/kg. Where multiple sediment contaminants exist, the Probable Effect Concentrations Quotient shall not exceed 0.75. See Appendix D and Section II, D for more information on the Probable Effect Concentrations Quotient.</p>

**Appendix C**

**Comment [R46]:** Previously Table 1.2  
Footnotes moved into "Notes" column

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING  
PURPOSES: NARRATIVE CRITERIA BASED ON NUMERIC THRESHOLDS NOT CONTAINED IN STATE WATER  
QUALITY STANDARDS (10 CSR 20-7.031)**

BENEFICIAL USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>ii</sup>	Notes
Protection of Aquatic Life	Biological: Aquatic Macro-invertebrates sampled using DNR Protocol.	3-4	<p><b>Full:</b> For seven or fewer samples and following DNR wadeable streams macroinvertebrate sampling and evaluation protocols, 75% of the stream condition index scores must be 16 or greater. Fauna achieving these scores are considered to be very similar to regional reference streams. For greater than seven samples or for other sampling and evaluation protocols, results must be statistically similar to representative reference or control stream.</p> <p><b>Non-Attainment:</b> For seven or fewer samples and following DNR wadeable streams macroinvertebrate sampling and evaluation protocols, 75% of the stream condition index scores must be 14 or lower. Fauna achieving these scores are considered to be substantially different from regional reference streams. For more than seven samples or for other sampling and evaluation protocols, results must be statistically dissimilar to control or representative reference streams.</p>	<p><b>Data Type Note:</b> DNR invert protocol will not be used for assessment in the Mississippi Alluvial Basin (bootheel area) due to lack of reference streams for comparison.</p> <p><b>Data Type Note:</b> See <a href="#">Section II.D</a> for additional criteria used to assess biological data.</p> <p><b>Compliance with Water Quality Standards Note:</b> See <a href="#">Appendix E</a>. For test streams that are significantly smaller than bioreference streams where both bioreference streams and small control streams are used to assess the biological integrity of the test stream, the assessment of the data should display and take into account both types of control streams.</p>
Protection of Aquatic Life	Biological: MDC Fish Community (RAM) Protocol (Ozark)	3-4	<p><b>Full:</b> For seven or fewer samples and following MDC RAM fish community protocols, 75% of the fIBI scores must be 36 or greater. Fauna achieving these scores are considered to be very similar to regional reference streams. For greater</p>	<p><b>Data Type Note:</b> See <a href="#">Section II.D</a> for additional criteria used to assess biological data.</p> <p><b>Compliance with Water Quality Standards Note:</b> <b>MDC fIBI scores</b> are from "Biological Criteria for Streams and Fish Communities in Missouri" by Doisy et al. (2008). If habitat limitations (as measured by either the QCPH1 index or other</p>

**Comment [R48]:** Previously Appendix E

**Comment [R49]:** Previously Table D-1 and B-2

**Comment [R50]:** Previously Appendix E

**Appendix C**

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NARRATIVE CRITERIA BASED ON NUMERIC THRESHOLDS NOT CONTAINED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

BENEFICIAL USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>ii</sup>	Notes
	Plateau only)		<p>than seven samples or for other sampling and evaluation protocols, results must be statistically similar to representative reference or control streams.</p> <p><u>Suspected of Impairment:</u> Data not conclusive (Category 2B or 3B). For first and second order streams IBI score &lt; 29.</p> <p><u>Non-Attainment:</u> First and second order streams will not be assessed for non-attainment. When assessing third to fifth order streams with data sets of seven or fewer samples collected by following MDC RAM fish community protocols, 75% of the IBI scores must be lower than 36. Fauna achieving these scores are considered to be substantially different from regional reference streams. For more than seven samples or for other sampling and evaluation protocols, results must be statistically dissimilar to control or representative reference streams.</p>	<p>appropriate methods) are judged to contribute to low fish community scores and this is the only type of data available, the water body will be included in Category 4C, 2B, or 3B. If other types of data exist, the weight of evidence approach will be used as described in this document.</p> <p><b>Compliance with Water Quality Standards Note:</b> For determining influence of poor habitat on those samples that are deemed as impaired, consultation with MDC RAM staff will be utilized. If, through this consultation, habitat is determined to be a significant possible cause for impairment, the water body will not be rated as impaired, but rather as suspect of impairment (categories 2B or 3B).</p> <p><b>Compliance with Water Quality Standards Note:</b> See Appendix D. For test streams that are significantly smaller than bioreference streams where both bioreference streams and small candidate reference streams are used to assess the biological integrity of the test stream, the assessment of the data should display and take into account both biocriteria reference streams and candidate reference streams.</p>
Protection of Aquatic Life	Other Biological Data	3-4	<p><u>Full:</u> Results must be statistically similar to representative reference or control streams.</p> <p><u>Non-Attainment:</u> Results must be statistically dissimilar to control or representative reference streams.</p>	<p><b>Data Type Note:</b> See Section II.D. for additional criteria used to assess biological data</p>

**Comment [R46]:** Previously Table 1.2  
Footnotes moved into "Notes" column

**Comment [R51]:** Previously Table B-1 and B-2

**Comment [R52]:** Previously Appendix E

**Appendix C**

**Comment [B46]:** Previously Table 1.2  
Footnotes moved into "Notes" column

**METHODS FOR ASSESSING COMPLIANCE WITH WATER QUALITY STANDARDS USED FOR 303(d) LISTING PURPOSES: NARRATIVE CRITERIA BASED ON NUMERIC THRESHOLDS NOT CONTAINED IN STATE WATER QUALITY STANDARDS (10 CSR 20-7.031)**

BENEFICIAL USES	DATA TYPE	DATA QUALITY CODE	COMPLIANCE WITH WATER QUALITY STANDARDS <sup>11</sup>	Notes
Protection of Aquatic Life	Toxicity testing of streams or lakes using aquatic organisms	2	<b>Full:</b> No more than one test result of statistically significant deviation from controls in acute or chronic test in a three-year period. <b>Non-Attainment:</b> Requirements for full attainment not met.	
<u>Human Health - Fish Consumption</u>	Chemicals (tissue)	1-2	<b>Full:</b> Contaminant levels in fish tissue levels in filets, tissue plugs, and eggs do not exceed guidelines. <b>Non-Attainment:</b> Requirements for full attainment not met.	<b>Compliance with Water Quality Standards Note:</b> Fish tissue threshold levels are: chlordane 0.1 mg/kg (Crellin, J.R. 1989, "New Trigger Levels for Chlordane in Fish-Revised Memo" Mo. Dept. of Health inter-office memorandum. June 16, 1989); mercury 0.3 mg/kg based on "Water Quality Criterion for Protection of Human Health: Methylmercury" EPA-823-R-01-001. Jan. 2001. <a href="http://www.epa.gov/waterscience/criteria/methylmercury/merctifl.pdf">http://www.epa.gov/waterscience/criteria/methylmercury/merctifl.pdf</a> ; PCBs 0.75 mg/kg, MDHSS Memorandum August 30, 2006 "Development of PCB Risk-based Fish Consumption Limit Tables;" and lead 0.3- mg/kg (World Health Organization 1972, "Evaluation of Certain Food Additives and the Contaminants Mercury, Lead and Cadmium." WHO Technical Report Series No. 505, Sixteenth Report on the Joint FAO/WHO Expert Committee on Food Additives. Geneva 33 pp. Assessment of Mercury will be based on samples solely from the following higher trophic level fish species: Walleye, Sauger, Trout, Black Bass, White Bass, Striped Bass, Northern Pike, Flathead Catfish and Blue Catfish. In a 2012 DHSS memorandum (not yet approved, but are being considered for future LMD revisions) threshold values are proposed to change as follows: chlordane 0.2 mg/kg; mercury 0.27 mg/kg; and PCBs = 0.540; lead has not changed, but they do add atrazine and PDBEs (Fish Fillet Advisory Concentrations (FFACs) in Missouri).

<sup>11</sup> See section on Statistical Considerations and Appendix D.

**Comment [B53]:** Previously Tables D-1 & B-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes
			Decision Rule/ Hypothesis	Criterion Used with the Decision Rule <sup>10</sup>	Significance Level ( $\alpha$ )	Decision Rule/ Hypothesis	Criterion Used with the Decision Rule	Significance Level ( $\alpha$ )	
Narrative Criteria	Color	Hypothesis Test: Two Sample, one tailed t-Test	Null Hypothesis: There is no difference in color between test stream and control stream.	Reject Null Hypothesis if calculated "t" value exceeds tabular "t" value for test alpha	0.1	Same Hypothesis	Same Criterion	0.1	
	Bottom deposits	Hypothesis Test, Two Sample, one tailed "t" Test	Null Hypothesis: Solids of anthropogenic origin cover less than 20% of stream bottom where velocity is less than 0.5 feet/second.	Reject Null Hypothesis if 60% Lower Confidence Limit (LCL) of mean percent fine sediment deposition (pfsd) in stream is greater than the sum of the pfsd in the control and 20 % more of the stream bottom. i.e., where the pfsd is expressed as a decimal, test stream pfsd > (control stream pfsd)+(0.20)	0.4	Same Hypothesis	Same Criterion	Same Significance Level	Criterion Note: If data is non-normal a nonparametric test will be used as a comparison of medians. The same 20% difference still applies. With current software the Mann- Whitney test is used.

Comment [D54]: Previously Table B-1

Comment [D55]: Previously Table B-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes						
			Decision Rule/ Hypothesis	Criterion Used with the Decision Rule <sup>ii</sup>	Significance Level ( <i>α</i> )	Decision Rule/ Hypothesis	Criteria Used with the Decision Rule	Significance Level ( <i>α</i> )							
Aquatic Life	Biological monitoring (Narrative)	For DNR Invert protocol: Sample sizes of 7 or less, 75% of samples must score 14 or lower.	Using DNR Invert Protocol: Null Hypothesis: Frequency of full sustaining scores for test stream is the same as for biological criteria reference streams.	Reject Null Hypothesis: if frequency of fully sustaining scores on test stream is significantly less than for biological criteria reference streams.	Not Applicable	Same Hypothesis	Same Criterion	Same Significance Level							
		For RAM Fish IBI protocol: Sample sizes of 7 or less, 75% of samples must score less than 36.													
		For DNR Invert protocol and sample size of 8 or more. Binomial Probability							A direct comparison of frequencies between test and biological criteria reference streams will be made.	Rate as impaired if biological criteria reference stream frequency of fully biologically supporting scores is greater than five percent more than test stream.	0.1	Same Hypothesis	Same Criterion	0.4	Criterion Note: For inverts, the reference number will change depending on which EDU the stream is in (X%-5%), for RAM samples the reference number will always be 70 (75%-5%).
		For RAM Fish IBI protocol and sample size of 8 or more. Binomial Probability.													
		For other biological data an appropriate parametric or							Null Hypothesis, Community metric(s) in	Reject Null Hypothesis if metric scores for test stream are	0.1	Same Hypothesis	Same Criterion	0.4	

Comment [284]: Primary Table B-1  
Comment [285]: Primary Table B-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes
			Decision Rule/ Hypothesis	Criterion Used with the Decision Rule <sup>(a)</sup>	Significance Level (a)	Decision Rule/ Hypothesis	Criterion Used with the Decision Rule	Significance Level (a)	
Aquatic Life (cont.)		nonparametric test will be used.	test stream is the same as for a reference stream or control streams.	significantly less than reference or control streams.					
			Other biological monitoring to be determined by type of data.	Dependent upon available information.	Dependent upon available information.	Same Hypothesis	Same Criterion	Same Significance Level	
	Toxic chemicals in water: (Numeric)	Not applicable	No more than one toxic event, toxicity test failure or exceedence of acute or chronic criterion in 3 years	Not applicable	Not applicable	Same Hypothesis	Same Criterion	Same Significance Level	
	Toxic chemicals in sediments: (Narrative)	Comparison of geometric mean to PEC value, or calculation of a PECQ value.	Waters are judged to be impaired if parameter geomean exceeds PEC, or site PECQ is exceeded.	For metals use 150% PEC threshold. The PECQ threshold value is 0.75.	Not applicable	Water is judged to be unimpaired if parameter geomean is equal to or less than PEC, or site PECQ equalled or not exceeded.	For metals use 150% of PEC threshold. The PECQ threshold value is 0.75.	Not applicable	<b>Compliance with Water Quality Standards</b> Note: In the case of toxic chemicals occurring in benthic sediment rather than in water, the numeric thresholds used to determine the need for further evaluation will be the Probable Effect Concentrations proposed in "Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems" by MacDonald, D.D. <i>et al.</i> Arch. Environ. Contam. Toxicol. 39,20-31 (2000). These Probable Effect Concentrations are as follows:

Comment [D54]: Previously Table B-1

Comment [D55]: Previously Table B-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes
			Decision Rule/ Hypothesis	Criteria Used with the Decision Rule <sup>ii</sup>	Significance Level ( <i>α</i> )	Decision Rule/ Hypothesis	Criteria Used with the Decision Rule	Significance Level ( <i>α</i> )	
Aquatic Life (cont.)									33 mg/kg As; 4.98 mg/kg Cd; 111 mg/kg Cr; 149 mg/kg Cu; 48.6 mg/kg Ni; 128 mg/kg Pb; 459 mg/kg Zn; 561 µg/kg naphthalene; 1170 µg/kg phenanthrene; 1520 µg/kg pyrene; 1050 µg/kg benzo(a)anthracene; 1290 µg/kg chrysene; 1450 µg/kg benzo(a)pyrene; 22,800 µg/kg total polycyclic aromatic hydrocarbons; 676 µg/kg total PCBs; <u>chlordan</u> 17.6 ug/kg; Sum DDE 31.3 ug/kg; <u>lindane</u> (gamma-BHC) 4.99 ug/kg. Where multiple sediment contaminants exist, the Probable Effect Concentrations Quotient shall not exceed 0.75. See Appendix D and Section II, D for more information on the Probable Effect Concentrations Quotient.
	Temperature, pH, total dissolved solids, oil and grease, dissolved oxygen (Numeric)	Binomial probability	Null Hypothesis: No more than 10% of samples exceed the water quality criterion.	Reject Null Hypothesis if the Type I error rate is less than 0.1.	Not applicable	Same Hypothesis	Same Criterion	Same Significance Level	
Losing Streams	<i>E.coli</i>	Binomial probability	Null Hypothesis: No more than 10% of samples exceed the water quality criterion.	Reject Null Hypothesis if the Type I error rate is less than 0.1.	0.1	Same Hypothesis	Same Criterion	Same Significance Level	

Comment [D54]: Previous Table 2-1  
Comment [D43]: Previous Table 2-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes
			Decision Rule/ Hypothesis	Criterion Used with the Decision Rule <sup>14</sup>	Significance Level ( $\alpha$ )	Decision Rule/ Hypothesis	Criterion Used with the Decision Rule	Significance Level ( $\alpha$ )	
Human Health – Fish Consumption	Toxic chemicals in water (Numeric)	Hypothesis test: 1-sided confidence limit	Null Hypothesis: Levels of contaminants in water do not exceed criterion.	Reject Null Hypothesis if the 60% LCL is greater than the criterion value.	0.5	Same Hypothesis	Reject Null Hypothesis if the 60% UCL is greater than the criterion value.	Same Significance Level	
	Toxic chemicals in tissue (Narrative)	Four or more samples: Hypothesis test 1-sided confidence limit	Null Hypothesis: Levels in fillet samples or fish eggs do not exceed criterion.	Reject Null Hypothesis if the 60% LCL is greater than the criterion value.	0.4	Same Hypothesis	Reject null hypothesis if the 60% UCL is greater than the criterion value.	Same Significance Level	
Drinking Water Supply (Raw)	Toxic chemicals (Numeric)	Hypothesis test: 1-sided confidence limit	Null Hypothesis: Levels of contaminants do not exceed criterion.	Reject Null Hypothesis if the 60% LCL is greater than the criterion value.	0.4	Same Hypothesis	Reject null hypothesis if the 60% UCL is greater than the criterion value.	Same Significance Level	
	Non-toxic chemicals (Numeric)	Hypothesis test: 1-sided confidence limit	Null Hypothesis: Levels of contaminants do not exceed criterion.	Reject Null Hypothesis: if the 60% LCL is greater than the criterion value.	0.4	Same Hypothesis	Reject null hypothesis if the 60% UCL is greater than the criterion value.	Same Significance Level	
Drinking Water Supply (Finished)	Toxic chemicals	Methods stipulated by Safe Drinking Water Act.	Methods stipulated by Safe Drinking Water Act.	Methods stipulated by Safe Drinking Water Act.	Methods stipulated by Safe Drinking Water Act.	Same Hypothesis	Same Criterion	Same Significance Level	

Comment [D54]: Previously Table B-1

Comment [D55]: Previously Table B-2

**Appendix D**

**DESCRIPTION OF ANALYTICAL TOOLS USED FOR DETERMINING THE STATUS OF MISSOURI WATERS (11" X 14" FOLD OUT)**

Designated Use	Analytes	Analytical Tool	Determining when waters are impaired			Determining when waters are no longer impaired			Notes
			Decision Rule/ Hypothesis	Criteria Used with the Decision Rule <sup>iii</sup>	Significance Level ( $\alpha$ )	Decision Rule/ Hypothesis	Criteria Used with the Decision Rule	Significance Level ( $\alpha$ )	
Whole Body Contact and Secondary	Bacteria (Numeric)	Geometric mean	Null Hypothesis: Levels of contaminants do not exceed criterion.	Reject Null Hypothesis: if the geometric mean is greater than the criterion value.	Not Applicable	Same Hypothesis	Same Criterion	Not applicable	
Irrigation & Livestock Water	Toxic chemicals (Numeric)	Hypothesis test 1-Sided confidence limit	Null Hypothesis: Levels of contaminants do not exceed criterion.	Reject Null Hypothesis if the 60% LCL is greater than the criterion value.	0.4	Same Hypothesis	Reject null hypothesis if the 60% UCL is greater than the criterion value.	Same Significance Level	
Protection of Aquatic Life	Nutrients in lakes (Numeric)	Hypothesis test	Null hypothesis: Criteria are not exceeded.	Reject Null Hypothesis if 60% LCL value is greater than criterion value.	0.4	Same Hypothesis	Same Criterion	Same Significance Level	<b>Hypothesis Test Note:</b> State nutrient criteria require at least four samples per year taken near the outflow point of the lake (or reservoir) between May 1 and August 31 for at least four different, not necessarily consecutive, years.

Comment [D54]: Previously Table B-4

Comment [D55]: Previously Table B-4

<sup>iii</sup> Where hypothesis testing is used for media other than fish tissue, for data sets with five samples or fewer, a 75 percent confidence interval around the appropriate central tendencies will be used to determine use attainment status. Use attainment will be determined as follows: (1) If the criterion value is above this interval (all values within the interval are in conformance with the criterion), rate as unimpaired; (2) If the criterion value falls within this interval, rate as unimpaired and place in Category 2B or 3B; (3) If the criterion value is below this interval (all values within the interval are not in conformance with the criterion), rate as impaired. For fish tissue, this procedure will be used with the following changes: (1) it will apply only to sample sizes of less than four and, (2) a 50% confidence interval will be used in place of the 75% confidence interval.



Date: November 18, 2015

Time: 9:00 am to 4:00 pm (adjourned early)

Meeting: Bioassessment Workgroup Meeting

Subject: Proposed 2018 Listing Methodology Meeting Notes

Attendees:

Trish Rielly, MoDNR	trish.rielly@dnr.mo.gov
Randy Sarver, MoDNR	randy.sarver@dnr.mo.gov
Dave Michaelson, MoDNR	dave.michaelson@dnr.mo.gov
Steve Walker, MoDNR	steve.walker@dnr.mo.gov
Sam McCord, MoDNR	samuel.mccord@dnr.mo.gov
Robert Voss, MoDNR	robert.voss@dnr.mo.gov
John Hoke, MoDNR	john.hoke@dnr.mo.gov
Bill Whipps, MoDNR	bill.whipps@dnr.mo.gov
Colleen Meredith, MoDNR	colleen.meredith@dnr.mo.gov
Liz Evans, Missouri Department of Health and Senior Services	elizabeth.evans@health.mo.gov
Matt Combes, Missouri Department of Conservation	matt.combes@mdc.mo.gov
Mike McKee, Missouri Department of Conservation	mike.mckee@mdc.mo.gov
Leslie Holloway, Missouri Farm Bureau	lholloway@mofb.com
Robert Brundage, Newman, Comley and Ruth	rbrundage@ncrpc.com
Trent Stober, HDR	trent.stober@hdrinc.com
David Carani, Geosyntec	dcarani@geosyntec.com

The meeting was scheduled to allow stakeholders the opportunity to provide comments or ask clarifying questions regarding the proposed 2018 Listing Methodology Document (LMD). Agenda items were requested from stakeholders prior to the meeting. A suggested agenda was provided, but it was stated that other topics were welcome for discussion. The proposed 2018 LMD discussions started in a format that allowed participants the opportunity to provide page-by-page comments. In doing so, other specific topics or concerns were raised and discussed. Below is a summary of the discussions.

**Overview of updates from the 2016 listing Methodology Document**

Trish Rielly provided an overview of the purpose of the meeting and the general updates that were made to the 2016 LMD. Overall, the majority of the updates related to reformatting the document by moving information previously provided in the appendix and blending the information with similar information provided in the main body of the document. Where possible, tables were merged and table footnotes (e.g., Tables B, C and D) were moved under the "Notes" column header. Other updates were more grammatical in nature.



### **Data Codes**

Questions were raised regarding a wording addition to Data Codes Two, Three and Four (page 17), and the reason for the additional wording “composite or plug” and “a minimum of one”. The additional wording was added to clarify what type of sample was actually being collected by the department. It was suggested to add clarification to explain what is involved in a quantitative biological monitoring study. For example, multiple data collection events for a quantitative macroinvertebrate study includes: monitoring of multiple stream reaches and habitat conditions, stream habitat survey, water quality sampling, and the collection of aquatic macroinvertebrate data from multiple seasons. Staff will add additional clarification to this section.

### **Weight of Evidence Approach**

It was noted the italicized sentence below was removed under the Weight of Evidence Approach (page 18). It was suggested that if the process described is being followed, the sentence should be retained. Staff explained that the sentence was inadvertently removed during the LMD update and merging of information. It was agreed that this wording should be maintained.

*For those analytes with numeric thresholds, the threshold values given in Table 1.2 (now Appendix C) will trigger a weight of evidence analysis to determine the existence or likelihood of a use impairment and the appropriateness of proposing a 303(d) listing based on narrative criteria.*

It was suggested that examples of other relevant environmental data be included with fish, aquatic macroinvertebrate scores, fish tissue, or toxicity testing of water or sediment. The addition of physical and chemical data should be included with biological and toxicity data to better understand toxicity assessments, in particular sediment toxicity. Suggested wording will be provided by stakeholder comment letter.

Regarding the assessment of sediment data, it was also suggested the department consider multiple lines of evidence or sediment toxicity tests instead of basing sediment toxicity thresholds on the probable effect concentration (PEC) values developed by McDonald, et al. Staff stated that further discussions may be warranted and that the professionals who developed the PEC thresholds should be present to participate in the discussion. Specific streams were discussed where stakeholders believed additional lines of evidence are necessary. Additional comments will be provided by stakeholder comment letter.

### **Aquatic Macroinvertebrate Community Data**

It was suggested that wording pertaining to “aquatic macroinvertebrate assessments” be retained, instead of using “aquatic biological assessments” to describe macroinvertebrate community health as a function of water quality and habitat (page 20). To ensure clarity, staff will add this wording back into the document.

Within the same paragraph, the term “usually” was also added in association with comparing test streams to reference streams within the same Ecological Drainage Unit (EDU). Clarification was requested by stakeholders and staff provided clarification regarding the addition of this word (page 20). Staff stated that there are instances when similar EDUs may be combined. When EDUs are combined, it



generally relates to cold water streams, and/or when there is only one reference stream within an ecological region of the state (e.g. Mississippi Alluvial Plains).

### **Assessing Small Streams**

There was significant discussion regarding how headwater streams and effluent dominated streams will be assessed against candidate reference streams. Several questions were raised regarding stream class sizes and how those class sizes are defined. Other questions were asked about the status of a current project being completed by the University of Missouri-Columbia (UMC) regarding the process/methods for establishing headwater reference streams (page 26). Stakeholders requested additional discussions on this topic. Once the UMC project is completed, the department plans to have the results of the work presented at an appropriate Water Protection Forum meeting.

Specific questions were raised on how land use-land cover statistics are calculated, and what types of barriers are considered as a migration barrier (page 27). Staff described how land use-land cover information is derived. Examples of migration barriers were provided, including how low water bridge crossings or culverts may operate as barriers. Staff stated that field surveys would be necessary to identify if a migration barrier was present on a stream or stream reach.

### **Appendix C**

It was suggested that the word "significantly" be removed (pages 52 and 53) from the "notes" column since it was not mentioned in the text on page 25. "Headwater" or "smaller than Wadeable/Perennial" were suggested alternative wording. Discussions continued regarding the definitions of stream classifications provided in the water quality standards. It was suggested that a reference to Table I of the water quality standards be included following the wording for bioreference streams. Staff agreed the removal and addition of specific wording could be made. A stakeholder stated they would also supply suggested wording via comment letter.

### **Assessing Macroinvertebrate Communities from Poor/Inadequate Habitat**

Discussion was requested on how macroinvertebrate samples would be assessed if MSCI scores are less than 16, and the habitat score less than 75 percent (page 23). Horse Creek was provided as an example for how a water body in this situation would be assessed.

It was suggested the percent habitat score from reference be added to the assessment worksheets. Staff agreed this additional information could be added to future biological assessment worksheets.

### **Age of Data**

There was considerable discussion on how the department considers the age of data when conducting assessments. In the past, it was agreed that the department would add a statement to the assessment worksheet providing a justification for using older data. Staff recalled this agreement and indicated a statement would be added to future assessment worksheets.

### **Mercury in Fish Tissue**

There is reference to a Missouri Department of Conservation survey within the LMD and fish tissue assessment worksheets; the status and availability of that MDC document were discussed. It was stated that this document was currently being revised but would be made available by the Missouri Department of Conservation.



Clarification was also provided regarding Department of Health and Senior Services, Department of Conservation, and EPA's values regarding mercury in fish tissue, and if they were consistent with EPA guidance. Staff stated the mercury in fish tissue assessments follow and are consistent with EPA guidance.

**Follow-up to the 2016 Listing Methodology Meeting Discussions held in February 2014**

Stakeholders asked if all the suggested updates were made to the 2016 LMD. Staff stated that all updates were made, with the exception of adding a statement to assessment worksheets when older data was used during an assessment cycle. Staff will include a statement, as needed, on future assessment worksheets.

**Assessment of pH with the clarification in water quality standards**

The current water quality standards rulemaking is proposing to clarify pH as a 4-day chronic criterion. Stakeholders were interested in whether this proposed revision would have an effect on assessing pH in the future. Staff presented USGS pH data by EDU to illustrate the current median pH values per EDU. It was suggested by stakeholders that staff look at the pH data by stream size, and time of day to see if that would have an effect. For continuous data, a method was presented to consider how to incorporate a 4-day chronic period and 10 percent exceedance rate. Discussions will continue on how to potentially handle discrete and sonde data at the December 1 public availability meeting. Staff and stakeholders will continue to look at continuous and discrete data sets, and how neighboring states assess pH.

Meeting adjourned early at approximately 12:30 pm after all stakeholder questions were answered.

The department will hold a second public availability session for the 2016 303(d) List and 2018 LMD on December 1, 2015. The public hearing for these documents is scheduled for January 6, 2016, and the public notice period will close on January 31, 2016.